

Generalized Vanishing Mean Oscillation Spaces Associated with Divergence Form Elliptic Operators

Renjin Jiang and Dachun Yang*

Abstract. Let L be a divergence form elliptic operator with complex bounded measurable coefficients, ω a positive concave function on $(0, \infty)$ of strictly critical lower type $p_\omega \in (0, 1]$ and $\rho(t) = t^{-1}/\omega^{-1}(t^{-1})$ for $t \in (0, \infty)$. In this paper, the authors introduce the generalized VMO spaces $\text{VMO}_{\rho, L}(\mathbb{R}^n)$ associated with L , and characterize them via tent spaces. As applications, the authors show that $(\text{VMO}_{\rho, L}(\mathbb{R}^n))^* = B_{\omega, L^*}(\mathbb{R}^n)$, where L^* denotes the adjoint operator of L in $L^2(\mathbb{R}^n)$ and $B_{\omega, L^*}(\mathbb{R}^n)$ the Banach completion of the Orlicz-Hardy space $H_{\omega, L^*}(\mathbb{R}^n)$. Notice that $\omega(t) = t^p$ for all $t \in (0, \infty)$ and $p \in (0, 1]$ is a typical example of positive concave functions satisfying the assumptions. In particular, when $p = 1$, then $\rho(t) \equiv 1$ and $(\text{VMO}_{1, L}(\mathbb{R}^n))^* = H_{L^*}^1(\mathbb{R}^n)$, where $H_{L^*}^1(\mathbb{R}^n)$ was the Hardy space introduced by Hofmann and Mayboroda.

1 Introduction

John and Nirenberg [22] introduced the space $\text{BMO}(\mathbb{R}^n)$, which is defined to be the space of all $f \in L^1_{\text{loc}}(\mathbb{R}^n)$ that satisfy

$$\|f\|_{\text{BMO}(\mathbb{R}^n)} \equiv \sup_{\text{ball } B \subset \mathbb{R}^n} \frac{1}{|B|} \int_B |f(x) - f_B| dx < \infty,$$

where and in what follows, $f_B \equiv \frac{1}{|B|} \int_B f(x) dx$. The space $\text{BMO}(\mathbb{R}^n)$ is proved to be the dual of the Hardy space $H^1(\mathbb{R}^n)$ by Fefferman and Stein in [13].

Sarason [24] introduced the space $\text{VMO}(\mathbb{R}^n)$, which is defined to be the space of all $f \in \text{BMO}(\mathbb{R}^n)$ that satisfy

$$\lim_{c \rightarrow 0} \sup_{B \subset \mathbb{R}^n, r_B \leq c} \frac{1}{|B|} \int_B |f(x) - f_B| dx = 0.$$

In order to represent $H^1(\mathbb{R}^n)$ as a dual space, Coifman and Weiss [9] introduced the space $\text{CMO}(\mathbb{R}^n)$ as the closure of continuous functions with compact supports in the $\text{BMO}(\mathbb{R}^n)$ norm and showed that $(\text{CMO}(\mathbb{R}^n))^* = H^1(\mathbb{R}^n)$. For more general cases, we refer to Janson [18] and Bourdaud [7].

2000 *Mathematics Subject Classification.* Primary 42B35; Secondary 42B30, 46E30.

Key words and phrases. divergence form elliptic operator, Gaffney estimate, Orlicz function, Orlicz-Hardy space, BMO, VMO, CMO, molecule, dual.

Dachun Yang is supported by the National Natural Science Foundation (Grant No. 10871025) of China.

* Corresponding author.

Let L be a linear operator in $L^2(\mathbb{R}^n)$ that generates an analytic semigroup $\{e^{-tL}\}_{t \geq 0}$ with kernels satisfying an upper bound of Poisson type. The Hardy space $H_L^1(\mathbb{R}^n)$ and the BMO space $\text{BMO}_L(\mathbb{R}^n)$ associated with L were defined and studied in [2, 11]. Duong and Yan [12] further proved that $(H_L^1(\mathbb{R}^n))^* = \text{BMO}_{L^*}(\mathbb{R}^n)$, where and in what follows, L^* denotes the adjoint operator of L in $L^2(\mathbb{R}^n)$. Moreover, recently, Deng et al in [10] introduced the space $\text{VMO}_L(\mathbb{R}^n)$, the space of vanishing mean oscillation associated with L , and proved that $(\text{VMO}_L(\mathbb{R}^n))^* = H_{L^*}^1(\mathbb{R}^n)$. Also, Auscher and Russ [5] studied the Hardy space H_L^1 on strongly Lipschitz domains associated with a divergence form elliptic operator L whose heat kernels have the Gaussian upper bounds and regularity, and Auscher, McIntosh and Russ [4] treated the Hardy space H^1 associated with the Hodge Laplacian on a Riemannian manifold with doubling measure.

Let A be an $n \times n$ matrix with entries $\{a_{j,k}\}_{j,k=1}^n \subset L^\infty(\mathbb{R}^n, \mathbb{C})$ satisfying the uniform ellipticity condition, namely, there exist constants $0 < \lambda_A \leq \Lambda_A < \infty$ such that for all $\xi, \zeta \in \mathbb{C}^n$ and almost every $x \in \mathbb{R}^n$,

$$(1.1) \quad \lambda_A |\xi|^2 \leq \text{Re} \langle A(x)\xi, \xi \rangle \quad \text{and} \quad |\langle A(x)\xi, \zeta \rangle| \leq \Lambda_A |\xi| |\zeta|.$$

Then the second order divergence form operator is given by

$$(1.2) \quad Lf \equiv \text{div}(A \nabla f),$$

interpreted in the weak sense via a sesquilinear form. It is well known that the kernel of the heat semigroup $\{e^{-tL}\}_{t > 0}$ lacks pointwise estimates in general.

From now on, in this paper, we always let L be as in (1.2) and L^* the adjoint operator of L in $L^2(\mathbb{R}^n)$. Recently, Hofmann and Mayboroda [16, 17] studied the Hardy space $H_L^1(\mathbb{R}^n)$ and its dual space $\text{BMO}_{L^*}(\mathbb{R}^n)$. Indeed, Hofmann and Mayboroda [16] first defined the Hardy space $H_L^1(\mathbb{R}^n)$ via its molecular decomposition, then established several maximal function characterizations of $H_L^1(\mathbb{R}^n)$, and in particular, showed that $(H_L^1(\mathbb{R}^n))^* = \text{BMO}_{L^*}(\mathbb{R}^n)$. These results were generalized in [20] to the Orlicz-Hardy space $H_{\omega,L}(\mathbb{R}^n)$ and its dual space $\text{BMO}_{\rho,L^*}(\mathbb{R}^n)$, which contain the Hardy spaces $H_L^p(\mathbb{R}^n)$ for all $p \in (0, 1]$, the space $\text{BMO}_{L^*}(\mathbb{R}^n)$ and the Lipschitz spaces $\text{Lip}_{L^*}(\frac{1}{p} - 1, \mathbb{R}^n)$ for all $p \in (0, 1)$ as special cases.

Let ω be a positive concave function on $(0, \infty)$ of strictly critical lower type $p_\omega \in (0, 1]$ and $\rho(t) = t^{-1}/\omega^{-1}(t^{-1})$ for $t \in (0, \infty)$. Recall that $\omega(t) = t^p$ for all $t \in (0, \infty)$ and $p \in (0, 1]$ is a typical example of such positive concave functions. Motivated by [9, 18, 10, 16, 20], in this paper, we introduce the generalized VMO spaces $\text{VMO}_{\rho,L}(\mathbb{R}^n)$ associated with L , and characterize them via tent spaces. Then, we prove that $(\text{VMO}_{\rho,L}(\mathbb{R}^n))^* = B_{\omega,L^*}(\mathbb{R}^n)$, where $B_{\omega,L^*}(\mathbb{R}^n)$ denotes the Banach completion of $H_{\omega,L^*}(\mathbb{R}^n)$. When $\omega(t) = t$ for all $t \in (0, \infty)$, we denote $\text{VMO}_{\rho,L}(\mathbb{R}^n)$ simply by $\text{VMO}_L(\mathbb{R}^n)$. In this case, our result reads as $(\text{VMO}_L(\mathbb{R}^n))^* = H_{L^*}^1(\mathbb{R}^n)$. Finally, we show that the space $\text{CMO}(\mathbb{R}^n)$ is a subspace of $\text{VMO}_L(\mathbb{R}^n)$, and if $n \geq 3$, then there exists an operator L as in (1.2), constructed in [1, 14], such that $\text{CMO}(\mathbb{R}^n) \subsetneq \text{VMO}_L(\mathbb{R}^n)$. Moreover, when $n = 1, 2$, the spaces $\text{CMO}(\mathbb{R}^n)$ and $\text{VMO}_L(\mathbb{R}^n)$ coincide with equivalent norms, which is pointed to us by the referee.

Precisely, this paper is organized as follows. In Section 2, we recall some known definitions and notation on the divergence form elliptic operator L and Orlicz functions ω considered in this paper.

In Section 3, we introduce the generalized VMO spaces $\text{VMO}_{\rho,L}(\mathbb{R}^n)$ associated with L , and tent spaces $T_{\omega,v}^\infty(\mathbb{R}_+^{n+1})$ and give some basic properties of these spaces. In particular, we characterize the space $\text{VMO}_{\rho,L}(\mathbb{R}^n)$ via $T_{\omega,v}^\infty(\mathbb{R}_+^{n+1})$; see Theorem 3.1 below.

In Section 4, we prove that $(\text{VMO}_{\rho,L}(\mathbb{R}^n))^* = B_{\omega,L^*}(\mathbb{R}^n)$, where $B_{\omega,L^*}(\mathbb{R}^n)$ denotes the Banach completion of $H_{\omega,L^*}(\mathbb{R}^n)$; see Theorem 4.2 below. In particular, we have $(\text{VMO}_L(\mathbb{R}^n))^* = H_{L^*}^1(\mathbb{R}^n)$. Finally, in Proposition 4.2 below, we show that the space $\text{CMO}(\mathbb{R}^n)$ is a subspace of $\text{VMO}_L(\mathbb{R}^n)$, and if $n \geq 3$, then there exists an operator L as in (1.2) such that $\text{CMO}(\mathbb{R}^n) \subsetneq \text{VMO}_L(\mathbb{R}^n)$; moreover, when $n = 1, 2$, the spaces $\text{CMO}(\mathbb{R}^n)$ and $\text{VMO}_L(\mathbb{R}^n)$ coincide with equivalent norms.

Finally, we make some conventions. Throughout the paper, we denote by C a positive constant which is independent of the main parameters, but it may vary from line to line. The symbol $X \lesssim Y$ means that there exists a positive constant C such that $X \leq CY$; $B \equiv B(z_B, r_B)$ denotes an open ball with center z_B and radius r_B and $CB(z_B, r_B) \equiv B(z_B, Cr_B)$. Moreover, in what follows, for each ball $B \subset \mathbb{R}^n$ and $j \in \mathbb{N}$, we set $U_0(B) \equiv B$ and $U_j(B) \equiv 2^j B \setminus 2^{j-1} B$. Set $\mathbb{N} \equiv \{1, 2, \dots\}$ and $\mathbb{Z}_+ \equiv \mathbb{N} \cup \{0\}$. For any subset E of $\mathbb{R}^n$, we denote by E^c the set $\mathbb{R}^n \setminus E$.

2 Preliminaries

In this section, we recall some notions and notation on divergence form elliptic operators, describe some basic assumptions on Orlicz functions and also present some basic properties on them.

2.1 Some notions on the divergence form elliptic operator L

A family $\{S_t\}_{t>0}$ of operators is said to satisfy the L^2 off-diagonal estimates, which are also called the Gaffney estimates (see [16]), if there exist positive constants c, C and β such that for arbitrary closed sets $E, F \subset \mathbb{R}^n$,

$$\|S_t f\|_{L^2(F)} \leq C e^{-\left(\frac{\text{dist}(E,F)^2}{ct}\right)^\beta} \|f\|_{L^2(E)}$$

for every $t > 0$ and every $f \in L^2(\mathbb{R}^n)$ supported in E . Here and in what follows, for any $p \in (0, \infty]$ and $E \subset \mathbb{R}^n$, $\|f\|_{L^p(E)} \equiv \|f\chi_E\|_{L^p(\mathbb{R}^n)}$; for any sets $E, F \subset \mathbb{R}^n$, $\text{dist}(E, F) \equiv \inf\{|x - y| : x \in E, y \in F\}$.

The following results were obtained in [3, 16, 15].

Lemma 2.1 ([15]). *If two families, $\{S_t\}_{t>0}$ and $\{T_t\}_{t>0}$, of operators satisfy the Gaffney estimates, then so does $\{S_t T_t\}_{t>0}$. Moreover, there exist positive constants c, C and β such that for arbitrary closed sets $E, F \subset \mathbb{R}^n$,*

$$\|S_s T_t f\|_{L^2(F)} \leq C e^{-\left(\frac{\text{dist}(E,F)^2}{c \max\{s,t\}}\right)^\beta} \|f\|_{L^2(E)}$$

for every $s, t > 0$ and every $f \in L^2(\mathbb{R}^n)$ supported in E .

Lemma 2.2 ([3, 15]). *The families*

$$(2.1) \quad \{e^{-tL}\}_{t>0}, \quad \{tLe^{-tL}\}_{t>0},$$

as well as

$$(2.2) \quad \{(I + tL)^{-1}\}_{t>0},$$

are bounded on $L^2(\mathbb{R}^n)$ uniformly in t and satisfy the Gaffney estimates with positive constants c and C , depending on n, λ_A, Λ_A as in (1.1) only. For the operators in (2.1), $\beta = 1$; while in (2.2), $\beta = 1/2$.

Following [16], set $p_L \equiv \inf\{p \geq 1 : \sup_{t>0} \|e^{-tL}\|_{L^p(\mathbb{R}^n) \rightarrow L^p(\mathbb{R}^n)} < \infty\}$ and

$$\tilde{p}_L \equiv \sup \left\{ p \leq \infty : \sup_{t>0} \|e^{-tL}\|_{L^p(\mathbb{R}^n) \rightarrow L^p(\mathbb{R}^n)} < \infty \right\}.$$

It was proved by Auscher [1] that if $n = 1, 2$, then $p_L = 1$ and $\tilde{p}_L = \infty$, and if $n \geq 3$, then $p_L < 2n/(n+2)$ and $\tilde{p}_L > 2n/(n-2)$. Moreover, thanks to a counterexample given by Frehse [14], this range is sharp.

Lemma 2.3 ([16]). *Let $k \in \mathbb{N}$ and $p \in (p_L, \tilde{p}_L)$. Then the operator given by setting, for all $f \in L^p(\mathbb{R}^n)$ and $x \in \mathbb{R}^n$,*

$$\mathcal{S}_L^k f(x) \equiv \left(\iint_{\Gamma(x)} |(t^2 L)^k e^{-t^2 L} f(y)| \frac{dy dt}{t^{n+1}} \right)^{1/2},$$

is bounded on $L^p(\mathbb{R}^n)$.

2.2 Orlicz functions

Let ω be a positive function defined on $\mathbb{R}_+ \equiv (0, \infty)$. The function ω is said to be of upper type p (resp. lower type p) for certain $p \in [0, \infty)$, if there exists a positive constant C such that for all $t \geq 1$ (resp. $t \in (0, 1]$) and all $s \in (0, \infty)$,

$$(2.3) \quad \omega(st) \leq Ct^p \omega(s).$$

Obviously, if ω is of lower type p for certain $p > 0$, then $\lim_{t \rightarrow 0+} \omega(t) = 0$. So for the sake of convenience, if it is necessary, we may assume that $\omega(0) = 0$. If ω is of both upper type p_1 and lower type p_0 , then ω is said to be of type (p_0, p_1) . Let

$$p_\omega^+ \equiv \inf\{p > 0 : \text{there exists } C > 0 \text{ such that (2.3) holds for all } t \in [1, \infty), s \in (0, \infty)\},$$

and

$$p_\omega^- \equiv \sup\{p > 0 : \text{there exists } C > 0 \text{ such that (2.3) holds for all } t \in (0, 1], s \in (0, \infty)\}.$$

The function ω is said to be of strictly lower type p if for all $t \in (0, 1)$ and $s \in (0, \infty)$, $\omega(st) \leq t^p \omega(s)$, and for a such function ω , we define

$$p_\omega \equiv \sup\{p > 0 : \omega(st) \leq t^p \omega(s) \text{ holds for all } s \in (0, \infty) \text{ and } t \in (0, 1)\}.$$

It is easy to see that $p_\omega \leq p_\omega^- \leq p_\omega^+$ for all ω . In what follows, p_ω , p_ω^- and p_ω^+ are called the strictly critical lower type index, the critical lower type index and the critical upper type index of ω , respectively.

Remark 2.1. We claim that if p_ω is defined as above, then ω is also of strictly lower type p_ω . In other words, p_ω is attainable. In fact, if this is not the case, then there exist certain $s \in (0, \infty)$ and $t \in (0, 1)$ such that $\omega(st) > t^{p_\omega} \omega(s)$. Hence there exists $\epsilon \in (0, p_\omega)$ small enough such that $\omega(st) > t^{p_\omega - \epsilon} \omega(s)$, which is contrary to the definition of p_ω . Thus, ω is of strictly lower type p_ω .

We now introduce the following assumption.

Assumption (A). Let ω be a positive function defined on $\mathbb{R}_+$, which is of strictly lower type and its strictly lower type index $p_\omega \in (0, 1]$. Also assume that ω is continuous, strictly increasing and concave. $\square$

Notice that if ω satisfies Assumption (A), then $\omega(0) = 0$ and ω is obviously of upper type 1. Since ω is concave, it is subadditive. In fact, let $0 < s < t$, then

$$\omega(s+t) \leq \frac{s+t}{t} \omega(t) \leq \omega(t) + \frac{s}{t} \frac{t}{s} \omega(s) = \omega(s) + \omega(t).$$

For any concave function ω of strictly lower type p , if we set $\tilde{\omega}(t) \equiv \int_0^t \omega(s)/s ds$ for $t \in [0, \infty)$, then by [25, Proposition 3.1], $\tilde{\omega}$ is equivalent to ω , namely, there exists a positive constant C such that $C^{-1}\omega(t) \leq \tilde{\omega}(t) \leq C\omega(t)$ for all $t \in [0, \infty)$; moreover, $\tilde{\omega}$ is strictly increasing, concave, subadditive and continuous function of strictly lower type p . Since all our results are invariant on equivalent functions, we always assume that ω satisfies Assumption (A); otherwise, we may replace ω by $\tilde{\omega}$.

For example, if $\omega(t) = t^p$ with $p \in (0, 1]$, then $p_\omega = p_\omega^+ = p$; if $\omega(t) = t^{1/2} \ln(e^4 + t)$, then $p_\omega = p_\omega^+ = 1/2$.

Let ω satisfy Assumption (A). A measurable function f on $\mathbb{R}^n$ is said to be in the Lebesgue type space $L(\omega)$ if

$$\int_{\mathbb{R}^n} \omega(|f(x)|) dx < \infty.$$

Moreover, for any $f \in L(\omega)$, define

$$\|f\|_{L(\omega)} \equiv \inf \left\{ \lambda > 0 : \int_{\mathbb{R}^n} \omega \left(\frac{|f(x)|}{\lambda} \right) dx \leq 1 \right\}.$$

Since ω is strictly increasing, we define the function $\rho(t)$ on $\mathbb{R}_+$ by setting, for all $t \in (0, \infty)$,

$$(2.4) \quad \rho(t) \equiv \frac{t^{-1}}{\omega^{-1}(t^{-1})},$$

where and in what follows, ω^{-1} denotes the inverse function of ω . Then the types of ω and ρ have the following relation; see [25] for its proof.

Proposition 2.1. *Let $0 < p_0 \leq p_1 \leq 1$ and ω be an increasing function. Then ω is of type (p_0, p_1) if and only if ρ is of type $(p_1^{-1} - 1, p_0^{-1} - 1)$.*

Throughout the whole paper, we always assume that ω satisfies Assumption (A) and ρ is as in (2.4).

3 Spaces $\text{VMO}_{\rho,L}(\mathbb{R}^n)$ associated with L

In this section, we introduce the generalized vanishing mean oscillation spaces associated with L . We begin with some notions and notation.

Let $q \in (p_L, \tilde{p}_L)$, $M \in \mathbb{N}$ and $\epsilon \in (0, \infty)$. A function $\alpha \in L^q(\mathbb{R}^n)$ is called an $(\omega, q, M, \epsilon)_L$ -molecule adapted to B if there exists a ball B such that

- (i) $\|\alpha\|_{L^q(U_j(B))} \leq 2^{-j\epsilon} |2^j B|^{1/q-1} \rho(|2^j B|)^{-1}$, $j \in \mathbb{Z}_+$;
- (ii) For every $k = 1, \dots, M$ and $j \in \mathbb{Z}_+$, there holds

$$\|(r_B^{-2} L^{-1})^k \alpha\|_{L^q(U_j(B))} \leq 2^{-j\epsilon} |2^j B|^{1/q-1} \rho(|2^j B|)^{-1}.$$

Let ϵ and M be as above. We also introduce the space

$$(3.1) \quad \mathcal{M}_\omega^{M,\epsilon}(L) \equiv \{\mu \in L^2(\mathbb{R}^n) : \|\mu\|_{\mathcal{M}_\omega^{M,\epsilon}(L)} < \infty\},$$

where

$$\|\mu\|_{\mathcal{M}_\omega^{M,\epsilon}(L)} \equiv \sup_{j \geq 0} \left\{ 2^{j\epsilon} |B(0, 2^j)|^{1/2} \rho(|B(0, 2^j)|) \sum_{k=0}^M \|L^{-k} \mu\|_{L^2(U_j(B(0,1)))} \right\}.$$

Notice that if $\phi \in \mathcal{M}_\omega^{M,\epsilon}(L)$ with norm 1, then ϕ is an $(\omega, 2, M, \epsilon)$ -molecule adapted to $B(0, 1)$. Conversely, if α is an $(\omega, 2, M, \epsilon)$ -molecule adapted to certain ball, then $\alpha \in \mathcal{M}_\omega^{M,\epsilon}(L)$.

Let A_t denote either $(I + t^2 L)^{-1}$ or $e^{-t^2 L}$ and $f \in (\mathcal{M}_\omega^{M,\epsilon}(L))^*$, the dual of $\mathcal{M}_\omega^{M,\epsilon}(L)$. We claim that $(I - A_t^*)^M f \in L_{\text{loc}}^2(\mathbb{R}^n)$ in the sense of distributions. In fact, for any ball B , if $\psi \in L^2(B)$, then it follows from the Gaffney estimates via Lemmas 2.1 and 2.2 that $(I - A_t)^M \psi \in \mathcal{M}_\omega^{M,\epsilon}(L)$ for all $\epsilon > 0$ and any fixed $t \in (0, \infty)$. Thus,

$$|\langle (I - A_t^*)^M f, \psi \rangle| \equiv |\langle f, (I - A_t)^M \psi \rangle| \leq C(t, r_B, \text{dist}(B, 0)) \|f\|_{(\mathcal{M}_\omega^{M,\epsilon}(L))^*} \|\psi\|_{L^2(B)},$$

which implies that $(I - A_t^*)^M f \in L_{\text{loc}}^2(\mathbb{R}^n)$ in the sense of distributions.

Finally, for any $M \in \mathbb{N}$, define

$$(3.2) \quad \mathcal{M}_{\omega,L^*}^M(\mathbb{R}^n) \equiv \bigcap_{\epsilon > n(1/p_\omega - 1/p_\omega^+)} (\mathcal{M}_\omega^{M,\epsilon}(L))^*.$$

Definition 3.1. Let $q \in (p_L, \tilde{p}_L)$ and $M > \frac{n}{2}(\frac{1}{p_\omega} - \frac{1}{2})$. An element $f \in \mathcal{M}_{\omega,L}^M(\mathbb{R}^n)$ is said to be in $\text{BMO}_{\rho,L}^{q,M}(\mathbb{R}^n)$ if

$$\|f\|_{\text{BMO}_{\rho,L}^{q,M}(\mathbb{R}^n)} \equiv \sup_{B \subset \mathbb{R}^n} \frac{1}{\rho(|B|)} \left[\frac{1}{|B|} \int_B |(I - e^{-r_B^2 L})^M f(x)|^q dx \right]^{1/q} < \infty,$$

where the supremum is taken over all balls B of $\mathbb{R}^n$.

Remark 3.1. The spaces $\text{BMO}_{\rho,L}^{q,M}(\mathbb{R}^n)$ were introduced in [20]; moreover, if $\omega(t) = t$ for all $t \in (0, \infty)$, $\text{BMO}_{\rho,L}^{q,M}(\mathbb{R}^n)$ is the space $\text{BMO}_L(\mathbb{R}^n)$ introduced by Hofmann and Mayboroda [16]. Since the spaces $\text{BMO}_{\rho,L}^{q,M}(\mathbb{R}^n)$ coincide for all $q \in (p_L, \tilde{p}_L)$ and $M > \frac{n}{2}(\frac{1}{p_\omega} - \frac{1}{2})$ (see [20, Theorem 4.1]), in what follows, we denote $\text{BMO}_{\rho,L}^{q,M}(\mathbb{R}^n)$ simply by $\text{BMO}_{\rho,L}(\mathbb{R}^n)$.

Let us introduce a new space $\text{VMO}_{\rho,L}^M(\mathbb{R}^n)$ as a subspace of $\text{BMO}_{\rho,L}(\mathbb{R}^n)$.

Definition 3.2. Let $M > \frac{n}{2}(\frac{1}{p_\omega} - \frac{1}{2})$. An element $f \in \text{BMO}_{\rho,L}(\mathbb{R}^n)$ is said to be in $\text{VMO}_{\rho,L}^M(\mathbb{R}^n)$ if it satisfies the following limiting conditions $\gamma_1(f) = \gamma_2(f) = \gamma_3(f) = 0$, where

$$\gamma_1(f) \equiv \lim_{c \rightarrow 0} \sup_{B: r_B \leq c} \frac{1}{\rho(|B|)} \left(\frac{1}{|B|} \int_B |(I - e^{-r_B^2 L})^M f(x)|^2 dx \right)^{1/2},$$

$$\gamma_2(f) \equiv \lim_{c \rightarrow \infty} \sup_{B: r_B \geq c} \frac{1}{\rho(|B|)} \left(\frac{1}{|B|} \int_B |(I - e^{-r_B^2 L})^M f(x)|^2 dx \right)^{1/2},$$

and

$$\gamma_3(f) \equiv \lim_{c \rightarrow \infty} \sup_{B \subset B(0,c)^c} \frac{1}{\rho(|B|)} \left(\frac{1}{|B|} \int_B |(I - e^{-r_B^2 L})^M f(x)|^2 dx \right)^{1/2}.$$

For any $f \in \text{VMO}_{\rho,L}^M(\mathbb{R}^n)$, define $\|f\|_{\text{VMO}_{\rho,L}^M(\mathbb{R}^n)} \equiv \|f\|_{\text{BMO}_{\rho,L}(\mathbb{R}^n)}$.

We next present some properties of the space $\text{VMO}_{\rho,L}^M(\mathbb{R}^n)$. To this end, we first recall some notions of tent spaces; see [8].

Let $\Gamma(x) \equiv \{(y, t) \in \mathbb{R}_+^{n+1} : |x - y| < t\}$ denote the standard cone (of aperture 1) with vertex $x \in \mathbb{R}^n$. For any closed set F of $\mathbb{R}^n$, denote by $\mathcal{R}F$ the union of all cones with vertices in F , i.e., $\mathcal{R}F \equiv \cup_{x \in F} \Gamma(x)$; for any open set O in $\mathbb{R}^n$, denote the tent over O by $\hat{O}$, which is defined by $\hat{O} \equiv [\mathcal{R}(O^c)]^c$.

For any measurable function g on $\mathbb{R}_+^{n+1}$ and $x \in \mathbb{R}^n$, define

$$\mathcal{A}(g)(x) \equiv \left(\iint_{\Gamma(x)} |g(y, t)|^2 \frac{dy dt}{t^{n+1}} \right)^{1/2}$$

and

$$\mathcal{C}_\rho(g)(x) \equiv \sup_{\text{ball } B \ni x} \frac{1}{\rho(|B|)} \left(\frac{1}{|B|} \int_{\hat{B}} |g(y, t)|^2 \frac{dy dt}{t} \right)^{1/2}.$$

For $p \in (0, \infty)$, the tent space $T_2^p(\mathbb{R}_+^{n+1})$ is defined to be the space of all measurable functions g on $\mathbb{R}_+^{n+1}$ such that $\|g\|_{T_2^p(\mathbb{R}_+^{n+1})} \equiv \|\mathcal{A}(g)\|_{L^p(\mathbb{R}^n)} < \infty$. Let ω satisfy Assumption (A). The tent space $T_\omega(\mathbb{R}_+^{n+1})$ associated to the function ω is defined to be the space of measurable functions g on $\mathbb{R}_+^{n+1}$ such that $\mathcal{A}(g) \in L(\omega)$ with the norm defined by

$$\|g\|_{T_\omega(\mathbb{R}_+^{n+1})} \equiv \|\mathcal{A}(g)\|_{L(\omega)} = \inf \left\{ \lambda > 0 : \int_{\mathbb{R}^n} \omega \left(\frac{\mathcal{A}(g)(x)}{\lambda} \right) dx \leq 1 \right\};$$

the space $T_\omega^\infty(\mathbb{R}_+^{n+1})$ is defined to be the space of all measurable functions g on $\mathbb{R}_+^{n+1}$ satisfying $\|g\|_{T_\omega^\infty(\mathbb{R}_+^{n+1})} \equiv \|\mathcal{C}_\rho(g)\|_{L^\infty(\mathbb{R}^n)} < \infty$.

In what follows, let $T_{\omega,v}^\infty(\mathbb{R}_+^{n+1})$ be the space of all $f \in T_\omega^\infty(\mathbb{R}_+^{n+1})$ satisfying $\eta_1(f) = \eta_2(f) = \eta_3(f) = 0$, where

$$\begin{aligned} \eta_1(f) &\equiv \lim_{c \rightarrow 0} \sup_{B: r_B \leq c} \frac{1}{\rho(|B|)} \left(\frac{1}{|B|} \iint_{\widehat{B}} |f(y, t)|^2 \frac{dy dt}{t} \right)^{1/2}, \\ \eta_2(f) &\equiv \lim_{c \rightarrow \infty} \sup_{B: r_B \geq c} \frac{1}{\rho(|B|)} \left(\frac{1}{|B|} \iint_{\widehat{B}} |f(y, t)|^2 \frac{dy dt}{t} \right)^{1/2}, \end{aligned}$$

and

$$\eta_3(f) \equiv \lim_{c \rightarrow \infty} \sup_{B \subset B(0, c)^c} \frac{1}{\rho(|B|)} \left(\frac{1}{|B|} \iint_{\widehat{B}} |f(y, t)|^2 \frac{dy dt}{t} \right)^{1/2}.$$

It is easy to see that $T_{\omega,v}^\infty(\mathbb{R}_+^{n+1})$ is a closed linear subspace of $T_\omega^\infty(\mathbb{R}_+^{n+1})$.

Further, denote by $T_{\omega,1}^\infty(\mathbb{R}_+^{n+1})$ the space of all $f \in T_\omega^\infty(\mathbb{R}_+^{n+1})$ with $\eta_1(f) = 0$, and $T_{2,c}^2(\mathbb{R}_+^{n+1})$ the space of all $f \in T_2^2(\mathbb{R}_+^{n+1})$ with compact supports. Obviously, we have $T_{2,c}^2(\mathbb{R}_+^{n+1}) \subset T_{\omega,v}^\infty(\mathbb{R}_+^{n+1}) \subset T_{\omega,1}^\infty(\mathbb{R}_+^{n+1})$. Finally, denote by $T_{\omega,0}^\infty(\mathbb{R}_+^{n+1})$ the closure of $T_{2,c}^2(\mathbb{R}_+^{n+1})$ in the space $T_{\omega,1}^\infty(\mathbb{R}_+^{n+1})$.

By [19, Proposition 3.1], we have the following result; see also [10].

Lemma 3.1. *Let $T_{\omega,v}^\infty(\mathbb{R}_+^{n+1})$ and $T_{\omega,0}^\infty(\mathbb{R}_+^{n+1})$ be defined as above. Then $T_{\omega,v}^\infty(\mathbb{R}_+^{n+1}) = T_{\omega,0}^\infty(\mathbb{R}_+^{n+1})$.*

Recall that a measure $d\mu$ on $\mathbb{R}_+^{n+1}$ is called a ρ -Carleson measure if

$$\sup_{B \subset \mathbb{R}^n} \frac{1}{\rho(|B|)} \left\{ \frac{1}{|B|} \iint_{\widehat{B}} |d\mu| \right\}^{1/2} < \infty,$$

where the supremum is taken over all balls B of $\mathbb{R}^n$.

We now characterize the space $\text{VMO}_{\rho,L}^M(\mathbb{R}^n)$ via tent spaces.

Theorem 3.1. *Let $M, M_1 \in \mathbb{N}$ and $M_1 \geq M > \frac{n}{2}(\frac{1}{p_\omega} - \frac{1}{2})$. Then the following conditions are equivalent:*

- (a) $f \in \text{VMO}_{\rho,L}^M(\mathbb{R}^n)$;
- (b) $f \in \mathcal{M}_{\omega,L}^{M_1}(\mathbb{R}^n)$ and $(t^2 L)^{M_1} e^{-t^2 L} f \in T_{\omega,v}^\infty(\mathbb{R}_+^{n+1})$.

Moreover, $\|(t^2 L)^{M_1} e^{-t^2 L} f\|_{T_{\omega,v}^\infty(\mathbb{R}_+^{n+1})}$ is equivalent to $\|f\|_{\text{BMO}_{\rho,L}(\mathbb{R}^n)}$.

To prove Theorem 3.1, we need two auxiliary results.

Let $M \in \mathbb{Z}_+$. In what follows, let C_M be the positive constant such that

$$(3.3) \quad C_M \int_0^\infty t^{2(M+1)} e^{-2t^2} \frac{dt}{t} = 1.$$

The following lemma was established in [20, Proposition 4.6].

Lemma 3.2. *Let $\epsilon \in (0, \infty)$ and $M > \frac{n}{2}(\frac{1}{p_\omega} - \frac{1}{2})$. If $f \in \text{BMO}_{\rho,L}(\mathbb{R}^n)$, then for any $(\omega, 2, M, \epsilon)_{L^*}$ -molecule α , there holds*

$$\int_{\mathbb{R}^n} f(x) \alpha(x) dx = C_M \iint_{\mathbb{R}_+^{n+1}} (t^2 L)^M e^{-t^2 L} f(x) t^2 L^* e^{-t^2 L^*} \alpha(x) \frac{dx dt}{t}.$$

Definition 3.3. *Let $M > \frac{n}{2}(\frac{1}{p_\omega} - \frac{1}{2})$. Define the space $\widetilde{\text{VMO}}_{\rho,L}^M(\mathbb{R}^n)$ to be the space of all elements $f \in \text{BMO}_{\rho,L}(\mathbb{R}^n)$ that satisfy the limiting conditions $\tilde{\gamma}_1(f) = \tilde{\gamma}_2(f) = \tilde{\gamma}_3(f) = 0$, where*

$$\tilde{\gamma}_1(f) \equiv \lim_{c \rightarrow 0} \sup_{B: r_B \leq c} \frac{1}{\rho(|B|)} \left(\frac{1}{|B|} \int_B |(I - [I + r_B^2 L]^{-1})^M f(x)|^2 dx \right)^{1/2},$$

$$\tilde{\gamma}_2(f) \equiv \lim_{c \rightarrow \infty} \sup_{B: r_B \geq c} \frac{1}{\rho(|B|)} \left(\frac{1}{|B|} \int_B |(I - [I + r_B^2 L]^{-1})^M f(x)|^2 dx \right)^{1/2},$$

and

$$\tilde{\gamma}_3(f) \equiv \lim_{c \rightarrow \infty} \sup_{B \subset B(0,c)^c} \frac{1}{\rho(|B|)} \left(\frac{1}{|B|} \int_B |(I - [I + r_B^2 L]^{-1})^M f(x)|^2 dx \right)^{1/2}.$$

Proposition 3.1. *Let $p \in (0, 1]$ and $M > \frac{n}{2}(\frac{1}{p} - \frac{1}{2})$. Then $f \in \text{VMO}_{\rho,L}^M(\mathbb{R}^n)$ if and only if $f \in \widetilde{\text{VMO}}_{\rho,L}^M(\mathbb{R}^n)$*

Proof. Recall that it was proved in [20, Lemma 4.1] that

$$(3.4) \quad \|f\|_{\text{BMO}_{\rho,L}(\mathbb{R}^n)} \sim \sup_{B \subset \mathbb{R}^n} \frac{1}{\rho(|B|)} \left[\frac{1}{|B|} \int_B |(I - (I + r_B^2 L)^{-1})^M f(x)|^2 dx \right]^{1/2}.$$

Now suppose that $f \in \widetilde{\text{VMO}}_{\rho,L}^M(\mathbb{R}^n)$. To see $f \in \text{VMO}_{\rho,L}^M(\mathbb{R}^n)$, it suffices to show that

$$(3.5) \quad \frac{1}{\rho(|B|)|B|^{1/2}} \left(\int_B |(I - e^{-r_B^2 L})^M f(x)|^2 dx \right)^{1/2} \lesssim \sum_{k=0}^{\infty} 2^{-k} \delta_k(f, B),$$

where

$$(3.6) \quad \delta_k(f, B) \equiv \sup_{\{B' \subset 2^{k+1}B: r_{B'} \in [2^{-1}r_B, r_B]\}} \frac{1}{\rho(|B|)|B|^{1/2}}$$

$$\times \left\{ \int_{B'} |(I - [I + r_B^2 L]^{-1})^M f(x)|^2 dx \right\}^{1/2}.$$

Indeed, since $f \in \widetilde{\text{VMO}}_{\rho, L}^M(\mathbb{R}^n)$, by Definition 3.3 and (3.4), we have that $\delta_k(f, B) \lesssim \|f\|_{\text{BMO}_{\rho, L}(\mathbb{R}^n)}$ and for each $k \in \mathbb{N}$,

$$\lim_{c \rightarrow 0} \sup_{B: r_B \leq c} \delta_k(f, B) = \lim_{c \rightarrow \infty} \sup_{B: r_B \geq c} \delta_k(f, B) = \lim_{c \rightarrow \infty} \sup_{B \subset B(0, c)^c} \delta_k(f, B) = 0.$$

Then by the dominated convergence theorem for series, we have

$$\begin{aligned} \gamma_1(f) &= \lim_{c \rightarrow 0} \sup_{B: r_B \leq c} \frac{1}{\rho(|B|)|B|^{1/2}} \left(\int_B |(I - e^{-r_B^2 L})^M f(x)|^2 dx \right)^{1/2} \\ &\lesssim \sum_{k=1}^{\infty} 2^{-k} \lim_{c \rightarrow 0} \sup_{B: r_B \leq c} \delta_k(f, B) = 0. \end{aligned}$$

Similarly, we have that $\gamma_2(f) = \gamma_3(f) = 0$, and thus, $f \in \text{VMO}_{\rho, L}^M(\mathbb{R}^n)$.

Let us now prove (3.5). Write

$$(3.7) \quad f = (I - [I + r_B^2 L]^{-1})^M f + \{I - (I - [I + r_B^2 L]^{-1})^M\} f \equiv f_1 + f_2,$$

where by Lemma 2.2, we have

$$\begin{aligned} (3.8) \quad &\|(I - e^{-r_B^2 L})^M f_1\|_{L^2(B)} \\ &\leq \sum_{k=0}^{\infty} \|(I - e^{-r_B^2 L})^M (f_1 \chi_{U_k(B)})\|_{L^2(B)} \lesssim \sum_{k=0}^{\infty} e^{-c2^{2k}} \|f_1 \chi_{U_k(B)}\|_{L^2(\mathbb{R}^n)} \\ &\lesssim \rho(|B|)|B|^{1/2} \sum_{k=0}^{\infty} e^{-c2^{2k}} 2^{kn} \delta_k(f, B) \lesssim \rho(|B|)|B|^{1/2} \sum_{k=0}^{\infty} 2^{-k} \delta_k(f, B), \end{aligned}$$

where c is a positive constant and the third inequality follows from the fact that there exists a collection, $\{B_{k,1}, B_{k,2}, \dots, B_{k,N_k}\}$, of balls such that each ball $B_{k,i}$ is of radius r_B , $B(x_B, 2^{k+1}r_B) \subset \cup_{i=1}^{N_k} B_{k,i}$ and $N_k \lesssim 2^{nk}$.

To estimate the remaining term, by the formula that

$$(3.9) \quad I - (I - [I + r_B^2 L]^{-1})^M = \sum_{j=1}^M \frac{M!}{j!(M-j)!} (r_B^2 L)^{-j} (I - [I + r_B^2 L]^{-1})^M$$

(which relies on the fact that $(I - (I + r^2 L)^{-1})(r^2 L)^{-1} = (I + r^2 L)^{-1}$ for all $r \in (0, \infty)$), and the Minkowski inequality, we obtain

$$\begin{aligned} (3.10) \quad &\|(I - e^{-r_B^2 L})^M f_2\|_{L^2(B)} \\ &\lesssim \sum_{j=1}^M \left(\int_B \left| (I - e^{-r_B^2 L})^{M-j} \left[- \int_0^{r_B} \frac{s}{r_B^2} e^{-s^2 L} ds \right]^j f_1(x) \right|^2 dx \right)^{1/2} \end{aligned}$$

$$\begin{aligned}
&\lesssim \sum_{j=1}^M \sum_{k=0}^{M-j} \int_0^{r_B} \cdots \int_0^{r_B} \frac{s_1}{r_B^2} \cdots \frac{s_j}{r_B^2} \|e^{-(kr_B^2 + s_1^2 + \cdots + s_j^2)L} f_1\|_{L^2(B)} ds_1 \cdots ds_j \\
&\lesssim \sum_{j=1}^M \sum_{k=0}^{M-j} \int_0^{r_B} \cdots \int_0^{r_B} \frac{s_1}{r_B^2} \cdots \frac{s_j}{r_B^2} \sum_{i=0}^{\infty} e^{-\frac{c(2^i r_B)^2}{kr_B^2 + s_1^2 + \cdots + s_j^2}} \|f_1 \chi_{U_i(B)}\|_{L^2(\mathbb{R}^n)} ds_1 \cdots ds_j \\
&\lesssim \rho(|B|)|B|^{1/2} \sum_{i=0}^{\infty} e^{-\frac{c2^{2i}}{M}} 2^{in} \delta_i(f, B) \lesssim \rho(|B|)|B|^{1/2} \sum_{i=0}^{\infty} 2^{-i} \delta_i(f, B),
\end{aligned}$$

where c is a positive constant and in the penultimate inequality, we used the fact that $\int_0^{r_B} \cdots \int_0^{r_B} \frac{s_1}{r_B^2} \cdots \frac{s_j}{r_B^2} ds_1 \cdots ds_j \sim 1$. Combining the estimates (3.8) and (3.10), we obtain

(3.5), which further implies that $\widetilde{\text{VMO}}_{\rho,L}^M(\mathbb{R}^n) \subset \text{VMO}_{\rho,L}^M(\mathbb{R}^n)$.

By borrowing some ideas from the proof of [16, Lemma 8.1], then similarly to the proof above, we have that $\text{VMO}_{\rho,L}^M(\mathbb{R}^n) \subset \widetilde{\text{VMO}}_{\rho,L}^M(\mathbb{R}^n)$. We omit the details here, which completes the proof of Proposition 3.1. $\square$

Proof of Theorem 3.1. We first show that (a) implies (b). Let $f \in \text{VMO}_{\rho,L}^M(\mathbb{R}^n)$. By [20, Theorem 6.1], we have that $(t^2 L)^{M_1} e^{-t^2 L} f \in T_{\omega}^{\infty}(\mathbb{R}_+^{n+1})$. To see $(t^2 L)^{M_1} e^{-t^2 L} f \in T_{\omega,v}^{\infty}(\mathbb{R}_+^{n+1})$, we claim that it suffices to show that for all balls B ,

$$(3.11) \quad \frac{1}{\rho(|B|)|B|^{1/2}} \left(\iint_{\widehat{B}} |(t^2 L)^{M_1} e^{-t^2 L} f(x)|^2 \frac{dx dt}{t} \right)^{1/2} \lesssim \sum_{k=0}^{\infty} 2^{-k} \delta_k(f, B),$$

where $\delta_k(f, B)$ is as in (3.6). In fact, since $f \in \text{VMO}_{\rho,L}^M(\mathbb{R}^n) = \widetilde{\text{VMO}}_{\rho,L}^M(\mathbb{R}^n)$, we have that for each $k \in \mathbb{N}$, $\delta_k(f, B) \lesssim \|f\|_{\text{BMO}_{\rho,L}(\mathbb{R}^n)}$ and

$$\lim_{c \rightarrow 0} \sup_{B: r_B \leq c} \delta_k(f, B) = \lim_{c \rightarrow \infty} \sup_{B: r_B \geq c} \delta_k(f, B) = \lim_{c \rightarrow \infty} \sup_{B \subset B(0, c)^c} \delta_k(f, B) = 0.$$

Then by the dominated convergence theorem for series, we have

$$\begin{aligned}
\eta_1(f) &= \lim_{c \rightarrow 0} \sup_{B: r_B \leq c} \frac{1}{\rho(|B|)|B|^{1/2}} \left(\iint_{\widehat{B}} |(t^2 L)^{M_1} e^{-t^2 L} f(x)|^2 \frac{dx dt}{t} \right)^{1/2} \\
&\lesssim \sum_{k=1}^{\infty} 2^{-k} \lim_{c \rightarrow 0} \sup_{B: r_B \leq c} \delta_k(f, B) = 0.
\end{aligned}$$

Similarly, we have $\eta_2(f) = \eta_3(f) = 0$, and thus, $(t^2 L)^{M_1} e^{-t^2 L} f \in T_{\omega,v}^{\infty}(\mathbb{R}_+^{n+1})$.

Let us prove (3.11). Write $f \equiv f_1 + f_2$ as in (3.7). Then by Lemmas 2.2 and 2.3, similarly to the estimate (3.8), we obtain

$$(3.12) \quad \left(\iint_{\widehat{B}} |(t^2 L)^{M_1} e^{-t^2 L} f_1(x)|^2 \frac{dx dt}{t} \right)^{1/2}$$

$$\begin{aligned}
&\leq \sum_{k=0}^{\infty} \left(\iint_{\widehat{B}} |(t^2 L)^{M_1} e^{-t^2 L} (f_1 \chi_{U_k(B)})(x)|^2 \frac{dx dt}{t} \right)^{1/2} \\
&\lesssim \|f_1\|_{L^2(4B)} + \sum_{k=3}^{\infty} \left(\int_0^{r_B} \exp \left\{ -\frac{(2^k r_B)^2}{ct^2} \right\} \frac{dt}{t} \right)^{1/2} \|f_1 \chi_{U_k(B)}\|_{L^2(\mathbb{R}^n)} \\
&\lesssim \|f_1\|_{L^2(4B)} + \sum_{k=3}^{\infty} \left\{ \int_0^{r_B} \left[\frac{t^2}{(2^k r_B)^2} \right]^{n+1} \frac{dt}{t} \right\}^{1/2} \|f_1 \chi_{U_k(B)}\|_{L^2(\mathbb{R}^n)} \\
&\lesssim \rho(|B|)|B|^{1/2} \sum_{k=0}^{\infty} 2^{-k} \delta_k(f, B),
\end{aligned}$$

where c is a positive constant. Applying (3.9), Lemma 2.2 and $M_1 \geq M$ to f_2 yields that

$$\begin{aligned}
(3.13) \quad &\left(\iint_{\widehat{B}} |(t^2 L)^{M_1} e^{-t^2 L} f_2(x)|^2 \frac{dx dt}{t} \right)^{1/2} \\
&\lesssim \sum_{j=1}^M \left(\iint_{\widehat{B}} |(t^2 L)^{M_1} e^{-t^2 L} (r_B^2 L)^{-j} f_1(x)|^2 \frac{dx dt}{t} \right)^{1/2} \\
&\lesssim \sum_{j=1}^M \sum_{k=0}^{\infty} \left(\int_{\widehat{B}} \left(\frac{t^2}{r_B^2} \right)^{2j} |(t^2 L)^{M_1-j} e^{-t^2 L} (f_1 \chi_{U_k(B)})(x)|^2 \frac{dx dt}{t} \right)^{1/2} \\
&\lesssim \sum_{j=1}^M \left\{ \sum_{k=0}^{\infty} \left[\int_0^{r_B} \left(\frac{t^2}{r_B^2} \right)^{2j} \frac{dt}{t} \right]^{1/2} \right. \\
&\quad \left. + \sum_{k=3}^{\infty} \left[\int_0^{r_B} \left(\frac{t^2}{r_B^2} \right)^{2j} \exp \left\{ -\frac{(2^k r_B)^2}{ct^2} \right\} \frac{dt}{t} \right]^{1/2} \right\} \|f_1 \chi_{U_k(B)}\|_{L^2(\mathbb{R}^n)} \\
&\lesssim \|f_1\|_{L^2(4B)} + \sum_{k=3}^{\infty} \left[\int_0^{r_B} \left(\frac{t^2}{(2^k r_B)^2} \right)^{n+1} \frac{dt}{t} \right]^{1/2} \|f_1 \chi_{U_k(B)}\|_{L^2(\mathbb{R}^n)} \\
&\lesssim \rho(|B|)|B|^{1/2} \sum_{k=0}^{\infty} 2^{-k} \delta_k(f, B).
\end{aligned}$$

The estimates (3.12) and (3.13) imply (3.11), and hence, completes the proof that (a) implies (b).

Conversely, let $f \in \mathcal{M}_{\omega, L}^{M_1}(\mathbb{R}^n)$ and $(t^2 L)^{M_1} e^{-t^2 L} f \in T_{\omega, v}^{\infty}(\mathbb{R}_+^{n+1})$. By [20, Theorem 6.1], we have that $f \in \text{BMO}_{\rho, L}(\mathbb{R}^n)$. For any ball B , write

$$\begin{aligned}
\left(\int_B |(I - e^{-r_B^2 L})^M f(x)|^2 dx \right)^{1/2} &= \sup_{\|g\|_{L^2(B)} \leq 1} \left| \int_B (I - e^{-r_B^2 L})^M f(x) g(x) dx \right| \\
&= \sup_{\|g\|_{L^2(B)} \leq 1} \left| \int_{\mathbb{R}^n} f(x) (I - e^{-r_B^2 L^*})^M g(x) dx \right|.
\end{aligned}$$

Notice that for any $g \in L^2(B)$, $(I - e^{-r_B^2 L^*})^M g$ is a multiple of an $(\omega, 2, M, \epsilon)_{L^*}$ -molecule; see [16, p. 43] and [20]. Then by Lemma 3.2 and the Hölder inequality, we obtain

$$\begin{aligned}
& \left(\int_B |(I - e^{-r_B^2 L})^M f(x)|^2 dx \right)^{1/2} \\
&= \sup_{\|g\|_{L^2(B)} \leq 1} \left| C_{M_1} \iint_{\mathbb{R}_+^{n+1}} (t^2 L)^{M_1} e^{-t^2 L} f(x) t^2 L^* e^{-t^2 L^*} (I - e^{-r_B^2 L^*})^M g(x) \frac{dx dt}{t} \right| \\
&\lesssim \sum_{k=0}^{\infty} \left\{ \iint_{V_k(B)} |(t^2 L)^{M_1} e^{-t^2 L} f(x)|^2 \frac{dx dt}{t} \right\}^{1/2} \\
&\quad \times \sup_{\|g\|_{L^2(B)} \leq 1} \left\{ \iint_{V_k(B)} |(t^2 L^* e^{-t^2 L^*} (I - e^{-r_B^2 L^*})^M g(x)|^2 \frac{dx dt}{t} \right\}^{1/2} \equiv \sum_{k=0}^{\infty} \sigma_k(f, B) \mathbf{I}_k,
\end{aligned}$$

where $V_0(B) \equiv \widehat{B}$ and $V_k(B) \equiv (\widehat{2^k B}) \setminus (\widehat{2^{k-1} B})$ for $k \in \mathbb{N}$. In what follows, for $k \geq 2$, let $V_{k,1}(B) \equiv (\widehat{2^k B}) \setminus (2^{k-2} B \times (0, \infty))$ and $V_{k,2}(B) \equiv V_k(B) \setminus V_{k,1}(B)$.

For $k = 0, 1, 2$, by Lemmas 2.2 and 2.3, we obtain

$$\begin{aligned}
\mathbf{I}_k &= \sup_{\|g\|_{L^2(B)} \leq 1} \left\{ \iint_{V_k(B)} |(t^2 L^* e^{-t^2 L^*} (I - e^{-r_B^2 L^*})^M g(x)|^2 \frac{dx dt}{t} \right\}^{1/2} \\
&\lesssim \sup_{\|g\|_{L^2(B)} \leq 1} \|(I - e^{-r_B^2 L^*})^M g\|_{L^2(\mathbb{R}^n)} \lesssim 1.
\end{aligned}$$

Now for $k \geq 3$, write

$$\begin{aligned}
\mathbf{I}_k &\lesssim \sup_{\|g\|_{L^2(B)} \leq 1} \left\{ \iint_{V_{k,1}(B)} |(t^2 L^* e^{-t^2 L^*} (I - e^{-r_B^2 L^*})^M g(x)|^2 \frac{dx dt}{t} \right\}^{1/2} \\
&\quad + \sup_{\|g\|_{L^2(B)} \leq 1} \left\{ \iint_{V_{k,2}(B)} \dots \right\}^{1/2} \equiv \mathbf{I}_{k,1} + \mathbf{I}_{k,2}.
\end{aligned}$$

Since for any $(y, t) \in V_{k,2}(B)$, $t \geq 2^{k-2} r_B$, by the Minkowski inequality and Lemma 2.2, we obtain

$$\begin{aligned}
\mathbf{I}_{k,2} &\lesssim \sup_{\|g\|_{L^2(B)} \leq 1} \left\{ \iint_{V_{k,2}(B)} \left| t^2 L^* e^{-t^2 L^*} \left[- \int_0^{r_B^2} L^* e^{-s L^*} ds \right]^M g(x) \right|^2 \frac{dx dt}{t} \right\}^{1/2} \\
&\lesssim \sup_{\|g\|_{L^2(B)} \leq 1} \int_0^{r_B^2} \dots \int_0^{r_B^2} \left\{ \iint_{V_{k,2}(B)} |t^2 (L^*)^{M+1} \right. \\
&\quad \left. \times e^{-(t^2 + s_1 + \dots + s_M) L^*} g(x)|^2 \frac{dx dt}{t} \right\}^{1/2} ds_1 \dots ds_M
\end{aligned}$$

$$\begin{aligned}
&\lesssim \sup_{\|g\|_{L^2(B)} \leq 1} \int_0^{r_B^2} \cdots \int_0^{r_B^2} \left\{ \int_{2^{k-2}r_B}^{2^k r_B} \frac{t^4 \|g\|_{L^2(B)}^2}{(t^2 + s_1 + \cdots + s_M)^{2(M+1)}} \frac{dt}{t} \right\}^{1/2} ds_1 \cdots ds_M \\
&\lesssim 2^{-2kM}.
\end{aligned}$$

Similarly, we have that $I_{k,1} \lesssim 2^{-2kM}$. Combining the above estimates and the fact that ρ is of upper type $1/p_\omega - 1$, we finally obtain

$$\begin{aligned}
\frac{1}{\rho(|B|)|B|^{1/2}} \left(\int_B |(I - e^{-r_B^2 L})^M f(x)|^2 dx \right)^{1/2} &\lesssim \sum_{k=0}^{\infty} 2^{-2kM} \frac{1}{\rho(|B|)|B|^{1/2}} \sigma_k(f, B) \\
&\lesssim \sum_{k=0}^{\infty} 2^{-k[2M - n(\frac{1}{p_\omega} - \frac{1}{2})]} \frac{\sigma_k(f, B)}{\rho(|2^k B|)|2^k B|^{1/2}}.
\end{aligned}$$

Since $(t^2 L)^{M_1} e^{-t^2 L} f \in T_{\omega, v}^\infty(\mathbb{R}_+^{n+1}) \subset T_\omega^\infty(\mathbb{R}_+^{n+1})$, by $M > \frac{n}{2}(\frac{1}{p_\omega} - \frac{1}{2})$ and the dominated convergence theorem for series, we have

$$\begin{aligned}
\gamma_1(f) &= \lim_{c \rightarrow 0} \sup_{B: r_B \leq c} \frac{1}{\rho(|B|)|B|^{1/2}} \left(\int_B |(I - e^{-r_B^2 L})^M f(x)|^2 dx \right)^{1/2} \\
&\lesssim \sum_{k=0}^{\infty} 2^{-k[2M - n(\frac{1}{p_\omega} - \frac{1}{2})]} \lim_{c \rightarrow 0} \sup_{B: r_B \leq c} \frac{\sigma_k(f, B)}{\rho(|2^k B|)|2^k B|^{1/2}} = 0.
\end{aligned}$$

Similarly, $\gamma_2(f) = \gamma_3(f) = 0$, which implies that $f \in \text{VMO}_{\rho, L}^M(\mathbb{R}^n)$, and hence, completes the proof of Theorem 3.1. $\square$

Remark 3.2. (i) It follows from Theorem 3.1 that for all $M \in \mathbb{N}$ and $M > \frac{n}{2}(\frac{1}{p_\omega} - \frac{1}{2})$, the spaces $\text{VMO}_{\rho, L}^M(\mathbb{R}^n)$ coincide with equivalent norms. Thus, in what follows, we denote $\text{VMO}_{\rho, L}^M(\mathbb{R}^n)$ simply by $\text{VMO}_{\rho, L}(\mathbb{R}^n)$; in particular, if $\omega(t) \equiv t$ for all $t \in (0, \infty)$, then $\rho(t) \equiv 1$ and we denote $\text{VMO}_{\rho, L}^M(\mathbb{R}^n)$ simply by $\text{VMO}_L(\mathbb{R}^n)$.

(ii) When A has real entries, or when the dimension $n = 1$ or $n = 2$ in the case of complex entries, the heat kernels always satisfy the Gaussian pointwise estimates (see [6]), in these cases, the space $\text{VMO}_L(\mathbb{R}^n)$ here coincides with the one introduced by Deng et al in [10].

4 Dual spaces of $\text{VMO}_{\rho, L}(\mathbb{R}^n)$

In this section, we identify the dual spaces of $\text{VMO}_{\rho, L}(\mathbb{R}^n)$. We begin with some notions and known facts on tent spaces.

Recall that a function a on $\mathbb{R}_+^{n+1}$ is called a $T_\omega(\mathbb{R}_+^{n+1})$ -atom if there exists a ball $B \subset \mathbb{R}^n$ such that $\text{supp } a \subset \widehat{B}$ and $(\iint_{\widehat{B}} |a(x, t)|^2 \frac{dx dt}{t})^{1/2} \leq \frac{1}{\rho(|B|)|B|^{1/2}}$.

Definition 4.1. Let $p \in (0, 1)$. The space $\widetilde{T}_\omega(\mathbb{R}_+^{n+1})$ is defined to be the space of all $f = \sum_{j \in \mathbb{N}} \lambda_j a_j$ in $(T_\omega^\infty(\mathbb{R}_+^{n+1}))^*$, where $\{a_j\}_{j \in \mathbb{N}}$ are $T_\omega(\mathbb{R}_+^{n+1})$ -atoms and $\{\lambda_j\}_{j \in \mathbb{N}} \in \ell^1$. If $f \in \widetilde{T}_\omega(\mathbb{R}_+^{n+1})$, then define $\|f\|_{\widetilde{T}_\omega(\mathbb{R}_+^{n+1})} \equiv \inf \{ \sum_{j \in \mathbb{N}} |\lambda_j| \}$, where the infimum is taken over all the possible decompositions of f as above.

By [16, Lemma 3.1], $\tilde{T}_\omega(\mathbb{R}_+^{n+1})$ is a Banach space. Moreover, from Definition 4.1, it is easy to deduce that $T_\omega(\mathbb{R}_+^{n+1})$ is dense in $\tilde{T}_\omega(\mathbb{R}_+^{n+1})$; in other words, $\tilde{T}_\omega(\mathbb{R}_+^{n+1})$ is a Banach completion of $T_\omega(\mathbb{R}_+^{n+1})$. The following lemma is just [19, Lemma 4.1].

Lemma 4.1. $T_\omega(\mathbb{R}_+^{n+1}) \subset \tilde{T}_\omega(\mathbb{R}_+^{n+1})$, $T_\omega(\mathbb{R}_+^{n+1})$ is dense in $\tilde{T}_\omega(\mathbb{R}_+^{n+1})$ and there exists a positive constant C such that for all $f \in T_\omega(\mathbb{R}_+^{n+1})$, $\|f\|_{\tilde{T}_\omega(\mathbb{R}_+^{n+1})} \leq C\|f\|_{T_\omega(\mathbb{R}_+^{n+1})}$.

The following theorem was established in [26]; see also [19, Theorem 4.2].

Theorem 4.1. $(T_{\omega,v}^\infty(\mathbb{R}_+^{n+1}))^* = \tilde{T}_\omega(\mathbb{R}_+^{n+1})$.

Now, let us recall some notions on the Hardy spaces associated with L . For all $f \in L^2(\mathbb{R}^n)$ and $x \in \mathbb{R}^n$, define

$$\mathcal{S}_L f(x) \equiv \left(\iint_{\Gamma(x)} |t^2 L e^{-t^2 L} f(y)|^2 \frac{dy dt}{t^{n+1}} \right)^{1/2}.$$

The space $H_{\omega,L}(\mathbb{R}^n)$ is defined to be the completion of the set $\{f \in L^2(\mathbb{R}^n) : \mathcal{S}_L f \in L(\omega)\}$ with respect to the quasi-norm $\|f\|_{H_{\omega,L}(\mathbb{R}^n)} \equiv \|\mathcal{S}_L f\|_{L(\omega)}$.

The Orlicz-Hardy space $H_{\omega,L}(\mathbb{R}^n)$ was introduced and studied in [20]. If $\omega(t) \equiv t$ for all $t \in (0, \infty)$, then the space $H_{\omega,L}(\mathbb{R}^n)$ coincides with the Hardy space $H_L^1(\mathbb{R}^n)$, which was introduced and studied by Hofmann and Mayboroda [16] (see also [17]).

Definition 4.2. Let $M \in \mathbb{N}$, $M > \frac{n}{2}(\frac{1}{p_\omega} - \frac{1}{2})$ and $\epsilon \in (n(\frac{1}{p_\omega} - \frac{1}{p_\omega^+}), \infty)$. An element $f \in (\text{BMO}_{\rho,L^*}(\mathbb{R}^n))^*$ is said to be in the space $H_{\omega,L}^{M,\epsilon}(\mathbb{R}^n)$ if there exist $\{\lambda_j\}_{j=1}^\infty \subset \mathbb{C}$ and $(\omega, 2, M, \epsilon)_L$ -molecules $\{\alpha_j\}_{j=1}^\infty$ such that $f = \sum_{j=1}^\infty \lambda_j \alpha_j$ in $(\text{BMO}_{\rho,L^*}(\mathbb{R}^n))^*$ and

$$\Lambda(\{\lambda_j \alpha_j\}_j) \equiv \inf \left\{ \lambda > 0 : \sum_{j=1}^\infty |B_j|_\omega \left(\frac{|\lambda_j|}{\lambda |B_j|_\omega} \right) \leq 1 \right\} < \infty,$$

where for each j , α_j is adapted to the ball B_j .

If $f \in H_{\omega,L}^{M,\epsilon}(\mathbb{R}^n)$, then its norm is defined by $\|f\|_{H_{\omega,L}^{M,\epsilon}(\mathbb{R}^n)} \equiv \inf \Lambda(\{\lambda_j \alpha_j\}_j)$, where the infimum is taken over all the possible decompositions of f as above.

It was proved in [20, Theorem 5.1] that for all $M > \frac{n}{2}(\frac{1}{p_\omega} - \frac{1}{2})$ and $\epsilon \in (n(\frac{1}{p_\omega} - \frac{1}{p_\omega^+}), \infty)$, the spaces $H_{\omega,L}(\mathbb{R}^n)$ and $H_{\omega,L}^{M,\epsilon}(\mathbb{R}^n)$ coincide with equivalent norms.

Let us introduce the Banach completion of the space $H_L^p(\mathbb{R}^n)$.

Definition 4.3. Let $\epsilon \in (n(\frac{1}{p_\omega} - \frac{1}{p_\omega^+}), \infty)$ and $M > \frac{n}{2}(\frac{1}{p_\omega} - \frac{1}{2})$. The space $B_{\omega,L}^{M,\epsilon}(\mathbb{R}^n)$ is defined to be the space of all $f = \sum_{j \in \mathbb{N}} \lambda_j \alpha_j$ in $(\text{BMO}_{\rho,L^*}(\mathbb{R}^n))^*$, where $\{\lambda_j\}_{j \in \mathbb{N}} \in \ell^1$ and $\{\alpha_j\}_{j \in \mathbb{N}}$ are $(\omega, 2, M, \epsilon)_L$ -molecules. If $f \in B_{\omega,L}^{M,\epsilon}(\mathbb{R}^n)$, define $\|f\|_{B_{\omega,L}^{M,\epsilon}(\mathbb{R}^n)} \equiv \inf \{\sum_{j \in \mathbb{N}} |\lambda_j|\}$, where the infimum is taken over all the possible decompositions of f as above.

By [16, Lemma 3.1] again, we see that $B_{\omega,L}^{M,\epsilon}(\mathbb{R}^n)$ is a Banach space. Moreover, from Definition 4.2, it is easy to deduce that $H_{\omega,L}(\mathbb{R}^n)$ is dense in $B_{\omega,L}^{M,\epsilon}(\mathbb{R}^n)$. More precisely, we have the following lemma.

Lemma 4.2. *Let $\epsilon \in (n(\frac{1}{p_\omega} - \frac{1}{p_\omega^+}), \infty)$ and $M > \frac{n}{2}(\frac{1}{p_\omega} - \frac{1}{2})$. Then*

- (i) $H_{\omega,L}(\mathbb{R}^n) \subset B_{\omega,L}^{M,\epsilon}(\mathbb{R}^n)$ and the inclusion is continuous.
- (ii) For any $\epsilon_1 \in (n(\frac{1}{p_\omega} - \frac{1}{p_\omega^+}), \infty)$ and $M_1 > \frac{n}{2}(\frac{1}{p_\omega} - \frac{1}{2})$, the spaces $B_{\omega,L}^{M,\epsilon}(\mathbb{R}^n)$ and $B_{\omega,L}^{M_1,\epsilon_1}(\mathbb{R}^n)$ coincide with equivalent norms.

Proof. From Definition 4.3 and the molecular characterization of $H_{\omega,L}(\mathbb{R}^n)$, it is easy to deduce (i). Let us prove (ii). By symmetry, it suffices to show that $B_{\omega,L}^{M,\epsilon}(\mathbb{R}^n) \subset B_{\omega,L}^{M_1,\epsilon_1}(\mathbb{R}^n)$.

Let $f \in B_{\omega,L}^{M,\epsilon}(\mathbb{R}^n)$. By Definition 4.3, there exist $(\omega, 2, M, \epsilon)_L$ -molecules $\{\alpha_j\}_{j \in \mathbb{N}}$ and $\{\lambda_j\}_{j \in \mathbb{N}} \subset \mathbb{C}$ such that $f = \sum_{j \in \mathbb{N}} \lambda_j \alpha_j$ in $(\text{BMO}_{\rho,L^*}(\mathbb{R}^n))^*$ and $\sum_{j \in \mathbb{N}} |\lambda_j| \lesssim \|f\|_{B_{\omega,L}^{M,\epsilon}(\mathbb{R}^n)}$.

By (i), for each $j \in \mathbb{N}$, we have that $\alpha_j \in H_{\omega,L}(\mathbb{R}^n) \subset B_{\omega,L}^{M_1,\epsilon_1}(\mathbb{R}^n)$ and $\|\alpha_j\|_{B_{\omega,L}^{M_1,\epsilon_1}(\mathbb{R}^n)} \lesssim \|\alpha_j\|_{H_{\omega,L}(\mathbb{R}^n)} \lesssim 1$. Since $B_{\omega,L}^{M_1,\epsilon_1}(\mathbb{R}^n)$ is a Banach space, we obtain that $f \in B_{\omega,L}^{M_1,\epsilon_1}(\mathbb{R}^n)$ and $\|f\|_{B_{\omega,L}^{M_1,\epsilon_1}(\mathbb{R}^n)} \leq \sum_{j \in \mathbb{N}} |\lambda_j| \|\alpha_j\|_{B_{\omega,L}^{M_1,\epsilon_1}(\mathbb{R}^n)} \lesssim \|f\|_{B_{\omega,L}^{M,\epsilon}(\mathbb{R}^n)}$. Thus, $B_{\omega,L}^{M,\epsilon}(\mathbb{R}^n) \subset B_{\omega,L}^{M_1,\epsilon_1}(\mathbb{R}^n)$, which completes the proof of Lemma 4.2. $\square$

Since the spaces $B_{\omega,L}^{M,\epsilon}(\mathbb{R}^n)$ coincide for all $\epsilon \in (n(\frac{1}{p_\omega} - \frac{1}{p_\omega^+}), \infty)$ and $M > \frac{n}{2}(\frac{1}{p_\omega} - \frac{1}{2})$, in what follows, we denote $B_{\omega,L}^{M,\epsilon}(\mathbb{R}^n)$ simply by $B_{\omega,L}(\mathbb{R}^n)$.

Lemma 4.3. $(B_{\omega,L})^* = \text{BMO}_{\rho,L^*}(\mathbb{R}^n)$.

Proof. Since $(H_{\omega,L}(\mathbb{R}^n))^* = \text{BMO}_{\rho,L^*}(\mathbb{R}^n)$ and $H_{\omega,L}(\mathbb{R}^n) \subset B_{\omega,L}(\mathbb{R}^n)$, by duality, we have that $(B_{\omega,L}(\mathbb{R}^n))^* \subset \text{BMO}_{\rho,L^*}(\mathbb{R}^n)$.

Conversely, let $\epsilon \in (n(\frac{1}{p_\omega} - \frac{1}{p_\omega^+}), \infty)$, $M > \frac{n}{2}(\frac{1}{p_\omega} - \frac{1}{2})$ and $f \in \text{BMO}_{\rho,L^*}(\mathbb{R}^n)$. For any $g \in B_{\omega,L}(\mathbb{R}^n)$, by Definition 4.3, there exist $\{\lambda_j\}_{j \in \mathbb{N}} \subset \mathbb{C}$ and $(\omega, M, 2, \epsilon)_L$ -molecules $\{\alpha_j\}_{j \in \mathbb{N}}$ such that $g = \sum_{j \in \mathbb{N}} \lambda_j \alpha_j$ in $(\text{BMO}_{\rho,L^*}(\mathbb{R}^n))^*$ and $\sum_{j \in \mathbb{N}} |\lambda_j| \leq 2\|g\|_{B_{\omega,L}(\mathbb{R}^n)}$. Thus,

$$\begin{aligned} |\langle f, g \rangle| &\leq \sum_{j \in \mathbb{N}} |\lambda_j| |\langle f, \alpha_j \rangle| \lesssim \sum_{j \in \mathbb{N}} |\lambda_j| \|f\|_{\text{BMO}_{\rho,L^*}(\mathbb{R}^n)} \|\alpha_j\|_{H_{\omega,L}(\mathbb{R}^n)} \\ &\lesssim \|f\|_{\text{BMO}_{\rho,L^*}(\mathbb{R}^n)} \|g\|_{B_{\omega,L}(\mathbb{R}^n)}, \end{aligned}$$

which implies that $f \in (B_{\omega,L}(\mathbb{R}^n))^*$, and hence, completes the proof of Lemma 4.3. $\square$

Let $M \in \mathbb{N}$. For all $F \in L^2(\mathbb{R}_+^{n+1})$ with compact support, define

$$(4.1) \quad \pi_{L,M} F \equiv C_M \int_0^\infty (t^2 L)^M e^{-t^2 L} F(\cdot, t) \frac{dt}{t},$$

where C_M is the positive constant same as in (3.3).

Proposition 4.1. *Let $M \in \mathbb{N}$. Then the operator $\pi_{L,M}$, initially defined on $T_{2,c}^2(\mathbb{R}_+^{n+1})$, extends to a bounded linear operator*

- (i) *from $T_2^p(\mathbb{R}_+^{n+1})$ to $L^p(\mathbb{R}^n)$, if $p \in (p_L, \tilde{p}_L)$;*
- (ii) *from $T_\omega(\mathbb{R}_+^{n+1})$ to $H_{\omega,L}(\mathbb{R}^n)$, if $M > \frac{n}{2}(\frac{1}{p_\omega} - \frac{1}{2})$;*
- (iii) *from $\tilde{T}_\omega(\mathbb{R}_+^{n+1})$ to $B_{\omega,L}(\mathbb{R}^n)$, if $M > \frac{n}{2}(\frac{1}{p_\omega} - \frac{1}{2})$;*
- (iv) *from $T_{\omega,v}^\infty(\mathbb{R}_+^{n+1})$ to $\text{VMO}_{\rho,L}(\mathbb{R}^n)$.*

Proof. (i) and (ii) were established in [20]. Let us prove (iii).

By [19, Lemma 4.7], we have that $T_{2,c}^2(\mathbb{R}_+^{n+1})$ is dense in $\tilde{T}_\omega(\mathbb{R}_+^{n+1})$. Let $f \in T_{2,c}^2(\mathbb{R}_+^{n+1})$. By (ii) and Lemma 4.2, we obtain that $\pi_{L,M}f \in H_{\omega,L}(\mathbb{R}^n) \subset B_{\omega,L}(\mathbb{R}^n)$. Moreover, by Definition 4.1, there exist $T_\omega(\mathbb{R}_+^{n+1})$ -atoms $\{a_j\}_{j \in \mathbb{N}}$ and $\{\lambda_j\}_{j \in \mathbb{N}} \subset \mathbb{C}$ such that $f = \sum_{j \in \mathbb{N}} \lambda_j a_j$ in $(T_\omega^\infty(\mathbb{R}_+^{n+1}))^*$ and $\sum_j |\lambda_j| \lesssim \|f\|_{\tilde{T}_\omega(\mathbb{R}_+^{n+1})}$. In addition, for any $g \in \text{BMO}_{\rho,L^*}(\mathbb{R}^n)$, by [20, Theorem 6.1], we have that $(t^2 L^*)^M e^{-t^2 L^*} g \in T_\omega^\infty(\mathbb{R}_+^{n+1})$. Thus, by the fact that $(T_\omega(\mathbb{R}_+^{n+1}))^* = T_\omega^\infty(\mathbb{R}_+^{n+1})$ (see [21, Proposition 4.1]), we obtain

$$\begin{aligned} \langle \pi_{L,M}(f), g \rangle &= C_M \int_{\mathbb{R}_+^{n+1}} f(x, t) (t^2 L^*)^M e^{-t^2 L^*} g(x) \frac{dx dt}{t} \\ &= \sum_{j \in \mathbb{N}} \lambda_j C_M \int_{\mathbb{R}_+^{n+1}} a_j(x, t) (t^2 L^*)^M e^{-t^2 L^*} g(x) \frac{dx dt}{t} = \sum_{j \in \mathbb{N}} \lambda_j \langle \pi_{L,M}(a_j), g \rangle, \end{aligned}$$

which implies that $\pi_{L,M}(f) = \sum_{j \in \mathbb{N}} \lambda_j \pi_{L,M}(a_j)$ in $(\text{BMO}_{\rho,L^*}(\mathbb{R}^n))^*$. By (ii), we further obtain

$$\begin{aligned} \|\pi_{L,M}(f)\|_{B_{\omega,L}(\mathbb{R}^n)} &\leq \sum_{j \in \mathbb{N}} |\lambda_j| \|\pi_{L,M}(a_j)\|_{B_{\omega,L}(\mathbb{R}^n)} \\ &\lesssim \sum_j |\lambda_j| \|\pi_{L,M}(a_j)\|_{H_{\omega,L}(\mathbb{R}^n)} \lesssim \|f\|_{\tilde{T}_\omega(\mathbb{R}_+^{n+1})}. \end{aligned}$$

Since $T_{2,c}^2(\mathbb{R}_+^{n+1})$ is dense in $\tilde{T}_\omega(\mathbb{R}_+^{n+1})$, we see that $\pi_{L,M}$ extends to a bounded linear operator from $\tilde{T}_\omega(\mathbb{R}_+^{n+1})$ to $B_{\omega,L}(\mathbb{R}^n)$, which completes the proof of (iii).

Let us now prove (iv). By the definition of $T_{\omega,0}^\infty(\mathbb{R}_+^{n+1})$, we have that $T_{2,c}^2(\mathbb{R}_+^{n+1})$ is dense in $T_{\omega,0}^\infty(\mathbb{R}_+^{n+1})$. Then by Lemma 3.1, we obtain that $T_{2,c}^2(\mathbb{R}_+^{n+1})$ is dense in $T_{\omega,v}^\infty(\mathbb{R}_+^{n+1})$. Thus, to prove (iv), it suffices to show that $\pi_{L,M}$ maps $T_{2,c}^2(\mathbb{R}_+^{n+1})$ continuously into $\text{VMO}_{\rho,L}(\mathbb{R}^n)$.

Let $f \in T_{2,c}^2(\mathbb{R}_+^{n+1})$. By (i), we see that $\pi_{L,M}f \in L^2(\mathbb{R}^n)$. Notice that (3.1) and (3.2) with L and L^* exchanged implies that $L^2(\mathbb{R}^n) \subset \mathcal{M}_{\omega,L}^{M_1}(\mathbb{R}^n)$, where $M_1 \in \mathbb{N}$ and $M_1 > \frac{n}{2}(\frac{1}{p_\omega} - \frac{1}{2})$. Thus, $\pi_{L,M}f \in \mathcal{M}_{\omega,L}^{M_1}(\mathbb{R}^n)$. To show $\pi_{L,M}f \in \text{VMO}_{\rho,L}(\mathbb{R}^n)$, by Theorem 3.1, we only need to verify that $(t^2 L)^{M_1} e^{-t^2 L} \pi_{L,M}f \in T_{\omega,v}^\infty(\mathbb{R}_+^{n+1})$.

For any ball $B \equiv B(x_B, r_B)$, let $V_0(B) \equiv \widehat{B}$ and $V_k(B) \equiv (\widehat{2^k B}) \setminus (\widehat{2^{k-1} B})$ for any $k \in \mathbb{N}$. For all $k \in \mathbb{Z}_+$, let $f_k \equiv f \chi_{V_k(B)}$. Thus, for $k = 0, 1, 2$, by Lemma 2.3 and (i), we obtain

$$\left(\iint_{\widehat{B}} |(t^2 L)^{M_1} e^{-t^2 L} \pi_{L,M} f_k(x)|^2 \frac{dx dt}{t} \right)^{1/2} \lesssim \|\pi_{L,M} f_k\|_{L^2(\mathbb{R}^n)} \lesssim \|f_k\|_{T_2^2(\mathbb{R}_+^{n+1})}.$$

For $k \geq 3$, let $V_{k,1}(B) \equiv (\widehat{2^k B}) \setminus (2^{k-2}B \times (0, \infty))$ and $V_{k,2}(B) \equiv V_k(B) \setminus V_{k,1}(B)$. We further write $f_k = f_k \chi_{V_{k,1}(B)} + f_k \chi_{V_{k,2}(B)} \equiv f_{k,1} + f_{k,2}$. By the Minkowski inequality, Lemma 2.2 and the Hölder inequality, we obtain

$$\begin{aligned}
& \left(\iint_{\widehat{B}} |(t^2 L)^{M_1} e^{-t^2 L} \pi_{L,M} f_{k,2}(x)|^2 \frac{dx dt}{t} \right)^{1/2} \\
&= C_M \left(\iint_{\widehat{B}} \left| \int_{2^{k-2}r_B}^{2^k r_B} (t^2 L)^{M_1} e^{-t^2 L} (s^2 L)^M e^{-s^2 L} (f_{k,2}(\cdot, s))(x) \frac{ds}{s} \right|^2 \frac{dx dt}{t} \right)^{1/2} \\
&\lesssim \int_{2^{k-2}r_B}^{2^k r_B} \left(\iint_{\widehat{B}} \left| t^{2M_1} s^{2M} L^{M+M_1} e^{-(s^2+t^2)L} (f_{k,2}(\cdot, s))(x) \right|^2 \frac{dx dt}{t} \right)^{1/2} \frac{ds}{s} \\
&\lesssim \int_{2^{k-2}r_B}^{2^k r_B} \left(\int_0^{r_B} \left| \frac{t^{2M_1} s^{2M}}{(s^2+t^2)^{M+M_1}} \right|^2 \|f_{k,2}(\cdot, s)\|_{L^2(\mathbb{R}^n)}^2 \frac{dt}{t} \right)^{1/2} \frac{ds}{s} \\
&\lesssim 2^{-2kM_1} \int_{2^{k-2}r_B}^{2^k r_B} \|f_{k,2}(\cdot, s)\|_{L^2(\mathbb{R}^n)} \frac{ds}{s} \lesssim 2^{-2kM_1} \|f_{k,2}\|_{T_2^2(\mathbb{R}_+^{n+1})}.
\end{aligned}$$

Similarly, we have

$$\left(\iint_{\widehat{B}} |(t^2 L)^{M_1} e^{-t^2 L} \pi_{L,M} f_{k,1}(x)|^2 \frac{dx dt}{t} \right)^{1/2} \lesssim 2^{-2kM_1} \|f_{k,1}\|_{T_2^2(\mathbb{R}_+^{n+1})}.$$

Combining the above estimates, we finally obtain that

$$\begin{aligned}
& \frac{1}{\rho(|B|)|B|^{1/2}} \left(\iint_{\widehat{B}} |(t^2 L)^{M_1} e^{-t^2 L} \pi_{L,M} f(x)|^2 \frac{dx dt}{t} \right)^{1/2} \\
&\lesssim \sum_{k=0}^2 \frac{1}{\rho(|B|)|B|^{1/2}} \left(\iint_{\widehat{B}} |(t^2 L)^{M_1} e^{-t^2 L} \pi_{L,M} f_k(x)|^2 \frac{dx dt}{t} \right)^{1/2} \\
&\quad + \sum_{k=3}^{\infty} \sum_{i=1}^2 \frac{1}{\rho(|B|)|B|^{1/2}} \left(\iint_{\widehat{B}} |(t^2 L)^{M_1} e^{-t^2 L} \pi_{L,M} f_{k,i}(x)|^2 \frac{dx dt}{t} \right)^{1/2} \\
&\lesssim \sum_{k=0}^2 \frac{1}{\rho(|B|)|B|^{1/2}} \|f_k\|_{T_2^2(\mathbb{R}_+^{n+1})} + \sum_{k=3}^{\infty} \sum_{i=1,2} \frac{2^{-2kM_1}}{\rho(|B|)|B|^{1/2}} \|f_{k,i}\|_{T_2^2(\mathbb{R}_+^{n+1})} \\
&\lesssim \sum_{k=0}^{\infty} 2^{-k[2M_1 - n(\frac{1}{p_\omega} - 1/2)]} \frac{1}{\rho(|2^k B|)|2^k B|^{1/2}} \|f_k\|_{T_2^2(\mathbb{R}_+^{n+1})},
\end{aligned}$$

where $2M_1 > n(1/p_\omega - 1/2)$. Since $f \in T_{\omega,v}^\infty(\mathbb{R}_+^{n+1}) \subset T_\omega^\infty(\mathbb{R}_+^{n+1})$, we have

$$\frac{1}{\rho(|2^k B|)|2^k B|^{1/2}} \|f_k\|_{T_2^2(\mathbb{R}_+^{n+1})} \lesssim \|f\|_{T_\omega^\infty(\mathbb{R}_+^{n+1})},$$

and for all fixed $k \in \mathbb{N}$,

$$\lim_{c \rightarrow 0} \sup_{B: r_B \leq c} \frac{\|f_k\|_{T_2^2(\mathbb{R}_+^{n+1})}}{\rho(|2^k B|)|2^k B|^{1/2}} = \lim_{c \rightarrow \infty} \sup_{B: r_B \geq c} \frac{\|f_k\|_{T_2^2(\mathbb{R}_+^{n+1})}}{\rho(|2^k B|)|2^k B|^{1/2}}$$

$$= \lim_{c \rightarrow \infty} \sup_{B: B \subset (B(0,c))^c} \frac{\|f_k\|_{T_2^2(\mathbb{R}_+^{n+1})}}{\rho(|2^k B|)|2^k B|^{1/2}} = 0.$$

Thus, by Theorem 3.1, we have

$$\|\pi_{L,M} f\|_{\text{BMO}_{\rho,L}(\mathbb{R}^n)} \sim \|(t^2 L)^{M_1} e^{-t^2 L} \pi_{L,M} f\|_{T_\omega^\infty(\mathbb{R}_+^{n+1})} \lesssim \|f\|_{T_\omega^\infty(\mathbb{R}_+^{n+1})},$$

and by the dominated convergence theorem for series,

$$\begin{aligned} \eta_1((t^2 L)^{M_1} e^{-t^2 L} \pi_{L,M} f) &= \lim_{c \rightarrow 0} \sup_{B: r_B \leq c} \frac{1}{\rho(|B|)|B|^{1/2}} \left(\int_B |(t^2 L)^{M_1} e^{-t^2 L} \pi_{L,M} f(x)|^2 dx \right)^{1/2} \\ &\lesssim \sum_{k=0}^{\infty} 2^{-k[2M_1 - n(1/p - 1/2)]} \lim_{c \rightarrow 0} \sup_{B: r_B \leq c} \frac{\|f_k\|_{T_2^2(\mathbb{R}_+^{n+1})}}{\rho(|2^k B|)|2^k B|^{1/2}} = 0. \end{aligned}$$

Similarly, we have that $\eta_2((t^2 L)^{M_1} e^{-t^2 L} \pi_{L,M} f) = \eta_3((t^2 L)^{M_1} e^{-t^2 L} \pi_{L,M} f) = 0$, and thus, $(t^2 L)^{M_1} e^{-t^2 L} \pi_{L,M} f \in T_{\omega,v}^\infty(\mathbb{R}_+^{n+1})$, which completes the proof of Proposition 4.1. $\square$

Lemma 4.4. $\text{VMO}_{\rho,L}(\mathbb{R}^n) \cap L^2(\mathbb{R}^n)$ is dense in $\text{VMO}_{\rho,L}(\mathbb{R}^n)$.

Proof. Let $f \in \text{VMO}_{\rho,L}(\mathbb{R}^n)$ and $M > \frac{n}{2}(\frac{1}{p_\omega} - \frac{1}{2})$. Then Theorem 3.1 tells us that $h \equiv (t^2 L)^M e^{-t^2 L} f \in T_{\omega,v}^\infty(\mathbb{R}_+^{n+1})$. Similarly to the proof of Proposition 4.1, by Lemma 3.1, there exist $\{h_k\}_{k \in \mathbb{N}} \subset T_{2,c}^2(\mathbb{R}_+^{n+1}) \subset T_{\omega,v}^\infty(\mathbb{R}_+^{n+1})$ such that $\|h - h_k\|_{T_\omega^\infty(\mathbb{R}_+^{n+1})} \rightarrow 0$, as $k \rightarrow \infty$. Thus, by Proposition 4.1 (i) and (iv), we obtain that $\pi_{L,1} h_k \in L^2(\mathbb{R}^n) \cap \text{VMO}_{\rho,L}(\mathbb{R}^n)$ and

$$(4.2) \quad \|\pi_{L,1}(h - h_k)\|_{\text{BMO}_{\rho,L}(\mathbb{R}^n)} \lesssim \|h - h_k\|_{T_\omega^\infty(\mathbb{R}_+^{n+1})} \rightarrow 0,$$

as $k \rightarrow \infty$.

Let α be an $(\omega, 2, M, \epsilon)_{L^*}$ -molecule. Then by the definition of $H_{\omega,L^*}(\mathbb{R}^n)$, $e^{-t^2 L^*} \alpha \in T_\omega(\mathbb{R}_+^{n+1})$, which, together with Lemma 3.2, the facts that $(T_\omega(\mathbb{R}_+^{n+1}))^* = T_\omega^\infty(\mathbb{R}_+^{n+1})$ (see [21, Proposition 4.1]) and $(H_{\omega,L^*}(\mathbb{R}^n))^* = \text{BMO}_{\rho,L}(\mathbb{R}^n)$, further implies that

$$\begin{aligned} \int_{\mathbb{R}^n} f(x) \alpha(x) dx &= C_M \iint_{\mathbb{R}_+^{n+1}} (t^2 L)^M e^{-t^2 L} f(x) t^2 L^* e^{-t^2 L^*} \alpha(x) \frac{dx dt}{t} \\ &= \lim_{k \rightarrow \infty} C_M \iint_{\mathbb{R}_+^{n+1}} h_k(x, t) t^2 L^* e^{-t^2 L^*} \alpha(x) \frac{dx dt}{t} \\ &= \frac{C_M}{C_1} \lim_{k \rightarrow \infty} \int_{\mathbb{R}^n} (\pi_{L,1} h_k(x)) \alpha(x) dx = \frac{C_M}{C_1} \langle \pi_{L,1} h, \alpha \rangle. \end{aligned}$$

Since the set of finite combinations of molecules is dense in $H_{\omega,L^*}(\mathbb{R}^n)$, we then obtain that $f = \frac{C_M}{C_1} \pi_{L,1} h$ in $\text{BMO}_{\rho,L}(\mathbb{R}^n)$.

Now, for each $k \in \mathbb{N}$, let $f_k \equiv \frac{C_M}{C_1} \pi_{L,1} h_k$. Then $f_k \in \text{VMO}_{\rho,L}(\mathbb{R}^n) \cap L^2(\mathbb{R}^n)$; moreover, by (4.2), we have $\|f - f_k\|_{\text{BMO}_{\rho,L}(\mathbb{R}^n)} \rightarrow 0$, as $k \rightarrow \infty$, which completes the proof of Lemma 4.4. $\square$

In what follows, the symbol $\langle \cdot, \cdot \rangle$ in the following theorem means the duality between the space $\text{BMO}_{\rho,L}(\mathbb{R}^n)$ and the space $B_{\omega,L^*}(\mathbb{R}^n)$ in the sense of Lemma 4.3 with L and L^* exchanged. Let us now state the main theorem of this paper.

Theorem 4.2. *The dual space of $\text{VMO}_{\rho,L}(\mathbb{R}^n)$, $(\text{VMO}_{\rho,L}(\mathbb{R}^n))^*$, coincides with the space $B_{\omega,L^*}(\mathbb{R}^n)$ in the following sense:*

(i) *For any $g \in B_{\omega,L^*}(\mathbb{R}^n)$, define the linear functional ℓ by setting, for all $f \in \text{VMO}_{\rho,L}(\mathbb{R}^n)$,*

$$(4.3) \quad \ell(f) \equiv \langle f, g \rangle.$$

Then there exists a positive constant C independent of g such that $\|\ell\|_{(\text{VMO}_{\rho,L}(\mathbb{R}^n))^} \leq C\|g\|_{B_{\omega,L^*}(\mathbb{R}^n)}$.*

(ii) *Conversely, for any $\ell \in (\text{VMO}_{\rho,L}(\mathbb{R}^n))^*$, there exists $g \in B_{\omega,L^*}(\mathbb{R}^n)$, such that (4.3) holds and there exists a positive constant C independent of ℓ such that $\|g\|_{B_{\omega,L^*}(\mathbb{R}^n)} \leq C\|\ell\|_{(\text{VMO}_{\rho,L}(\mathbb{R}^n))^*}$.*

Proof. By Lemma 4.3, we have $(B_{\omega,L^*}(\mathbb{R}^n))^* = \text{BMO}_{\rho,L}(\mathbb{R}^n)$. By Definition 3.2, we see that $\text{VMO}_{\rho,L}(\mathbb{R}^n) \subset \text{BMO}_{\rho,L}(\mathbb{R}^n)$, which implies that $B_{\omega,L^*}(\mathbb{R}^n) \subset (\text{VMO}_{\rho,L}(\mathbb{R}^n))^*$.

Conversely, let $M > \frac{n}{2}(\frac{1}{p_\omega} - \frac{1}{2})$ and $\ell \in (\text{VMO}_{\rho,L}(\mathbb{R}^n))^*$. By Proposition 4.1, $\pi_{L,1}$ is bounded from $T_{\omega,v}^\infty(\mathbb{R}_+^{n+1})$ to $\text{VMO}_{\rho,L}(\mathbb{R}^n)$, which implies that $\ell \circ \pi_{L,1}$ is a bounded linear functional on $T_{\omega,v}^\infty(\mathbb{R}_+^{n+1})$. Thus, by Theorem 4.1, there exists $g \in \tilde{T}_\omega(\mathbb{R}_+^{n+1})$ such that for all $f \in T_{\omega,v}^\infty(\mathbb{R}_+^{n+1})$, $\ell \circ \pi_{L,1}(f) = \langle f, g \rangle$.

Now, suppose that $f \in \text{VMO}_{\rho,L}(\mathbb{R}^n) \cap L^2(\mathbb{R}^n)$. By Theorem 3.1, we have that $(t^2 L)^M e^{-t^2 L} f \in T_{\omega,v}^\infty(\mathbb{R}_+^{n+1})$. Moreover, by the proof of Lemma 4.4, we see that $f = \frac{C_M}{C_1} \pi_{L,1}((t^2 L)^M e^{-t^2 L} f)$ in $\text{BMO}_{\rho,L}(\mathbb{R}^n)$. Thus,

$$(4.4) \quad \ell(f) = \frac{C_M}{C_1} \ell \circ \pi_{L,1}((t^2 L)^M e^{-t^2 L} f) = \frac{C_M}{C_1} \iint_{\mathbb{R}_+^{n+1}} (t^2 L)^M e^{-t^2 L} f(x) g(x, t) \frac{dx dt}{t}.$$

By [19, Lemma 4.7], $T_{2,c}^2(\mathbb{R}_+^{n+1})$ is dense in $\tilde{T}_\omega(\mathbb{R}_+^{n+1})$. Since $g \in \tilde{T}_\omega(\mathbb{R}_+^{n+1})$, we may choose $\{g_k\}_{k \in \mathbb{N}} \subset T_{2,c}^2(\mathbb{R}_+^{n+1})$ such that $g_k \rightarrow g$ in $\tilde{T}_\omega(\mathbb{R}_+^{n+1})$. By Proposition 4.1 (iii), we have that $\pi_{L^*,M}(g)$, $\pi_{L^*,M}(g_k) \in B_{\omega,L^*}(\mathbb{R}^n)$ and

$$\|\pi_{L^*,M}(g - g_k)\|_{B_{\omega,L^*}(\mathbb{R}^n)} \lesssim \|g - g_k\|_{\tilde{T}_\omega(\mathbb{R}_+^{n+1})} \rightarrow 0,$$

as $k \rightarrow \infty$. This, together with (4.4), Theorem 4.1, the dominated convergence theorem and Lemma 4.3, implies that

$$(4.5) \quad \begin{aligned} \ell(f) &= \frac{C_M}{C_1} \lim_{k \rightarrow \infty} \iint_{\mathbb{R}_+^{n+1}} (t^2 L)^M e^{-t^2 L} f(x) g_k(x, t) \frac{dx dt}{t} \\ &= \frac{C_M}{C_1} \lim_{k \rightarrow \infty} \int_{\mathbb{R}^n} f(x) \int_0^\infty (t^2 L^*)^M e^{-t^2 L^*} (g_k(\cdot, t))(x) \frac{dt}{t} dx \\ &= \frac{1}{C_1} \lim_{k \rightarrow \infty} \langle f, \pi_{L^*,M}(g_k) \rangle = \frac{1}{C_1} \langle f, \pi_{L^*,M}(g) \rangle. \end{aligned}$$

Since $\text{VMO}_{\rho,L}(\mathbb{R}^n) \cap L^2(\mathbb{R}^n)$ is dense in $\text{VMO}_{\rho,L}(\mathbb{R}^n)$ (see Lemma 4.4), we finally obtain that (4.5) holds for all $f \in \text{VMO}_{\rho,L}(\mathbb{R}^n)$, and $\|\ell\|_{(\text{VMO}_{\rho,L}(\mathbb{R}^n))^*} = \frac{1}{C_1} \|\pi_{L^*,M}g\|_{B_{\omega,L^*}(\mathbb{R}^n)}$. In this sense, we have that $(\text{VMO}_{\rho,L}(\mathbb{R}^n))^* \subset B_{\omega,L^*}(\mathbb{R}^n)$, which completes the proof of Theorem 4.2. $\square$

Remark 4.1. If $\omega(t) \equiv t$ for all $t \in (0, \infty)$, then by Definitions 4.2 and 4.3, we have that $B_{\omega,L^*}(\mathbb{R}^n) = H_{\omega,L^*}(\mathbb{R}^n) = H_{L^*}^1(\mathbb{R}^n)$, where $H_{L^*}^1(\mathbb{R}^n)$ was the Hardy space introduced by Hofmann and Mayboroda in [16]. By Theorem 4.2, we obtain that $(\text{VMO}_{\rho,L}(\mathbb{R}^n))^* = (\text{VMO}_L(\mathbb{R}^n))^* = H_{L^*}^1(\mathbb{R}^n)$.

In what follows, let $\omega(t) \equiv t$ for all $t \in (0, \infty)$. We denote $(\omega, 2, M, \epsilon)_L$ -molecule simply by $(1, 2, M, \epsilon)_L$ -molecule.

We now compare the space $\text{VMO}_L(\mathbb{R}^n)$ with the classical space $\text{CMO}(\mathbb{R}^n)$ introduced by Coifman and Weiss [9].

Recall that the space $\text{BMO}(\mathbb{R}^n)$ is defined to be the set of all $f \in L_{\text{loc}}^1(\mathbb{R}^n)$ that satisfy

$$\|f\|_{\text{BMO}(\mathbb{R}^n)} \equiv \sup_{B \subset \mathbb{R}^n} \frac{1}{|B|} \int_B |f(x) - f_B| dx < \infty,$$

where $f_B \equiv \frac{1}{|B|} \int_B f(x) dx$. Then the space $\text{CMO}(\mathbb{R}^n)$ is defined to be the closure of $C_c(\mathbb{R}^n)$ (the set of all continuous functions with compact support) in the norm $\|\cdot\|_{\text{BMO}(\mathbb{R}^n)}$.

Proposition 4.2. *For all $n \in \mathbb{N}$, the space $\text{CMO}(\mathbb{R}^n)$ is continuously embedded in the space $\text{VMO}_L(\mathbb{R}^n)$. Moreover, when $n \geq 3$, then there exists L as in (1.2) such that $\text{CMO}(\mathbb{R}^n)$ is a proper subset of $\text{VMO}_L(\mathbb{R}^n)$.*

Proof. Let $M > \frac{n}{4}$ and $\epsilon \in (0, \infty)$. Notice that each $(1, 2, M, \epsilon)_L$ -molecule is a classical $H^1(\mathbb{R}^n)$ -molecule up to a harmless constant; see [16, p. 41] or [20, Remark 7.1]. Thus, we have that $H_{L^*}^1(\mathbb{R}^n) \subset H^1(\mathbb{R}^n)$ and for all $f \in H_{L^*}^1(\mathbb{R}^n)$, $\|f\|_{H^1(\mathbb{R}^n)} \lesssim \|f\|_{H_{L^*}^1(\mathbb{R}^n)}$.

Since $(H_{L^*}^1(\mathbb{R}^n))^* = \text{BMO}_L(\mathbb{R}^n)$ and $(H^1(\mathbb{R}^n))^* = \text{BMO}(\mathbb{R}^n)$, we further obtain that $\text{BMO}(\mathbb{R}^n) \subset \text{BMO}_L(\mathbb{R}^n)$, and for all $g \in \text{BMO}(\mathbb{R}^n)$, $\|g\|_{\text{BMO}_L(\mathbb{R}^n)} \lesssim \|g\|_{\text{BMO}(\mathbb{R}^n)}$.

Let us now show that $C_c(\mathbb{R}^n) \subset \text{VMO}_L(\mathbb{R}^n)$. Suppose that $f \in C_c(\mathbb{R}^n)$ and $B \equiv B(x_B, r_B)$. By an argument as in [1, Section 2.5], we have that for all $t > 0$, $e^{-tL}1 = 1$ in the $L_{\text{loc}}^2(\mathbb{R}^n)$ sense, that is, for any $\phi \in L^2(\mathbb{R}^n)$ with compact support, there holds

$$\int_{\mathbb{R}^n} \phi(x) dx = \int_{\mathbb{R}^n} e^{-tL} 1 \phi(x) dx = \int_{\mathbb{R}^n} e^{-tL^*} \phi(x) dx.$$

This together with Lemma 2.2 and the Hölder inequality implies that

$$\begin{aligned} \|(I - e^{-r_B^2 L})^M f\|_{L^2(B)} &= \sup_{\|g\|_{L^2(B)} \leq 1} \left| \int_B (I - e^{-r_B^2 L})^M f(x) g(x) dx \right| \\ &= \sup_{\|g\|_{L^2(B)} \leq 1} \left| \int_B (I - e^{-r_B^2 L})^M (f - f_B)(x) g(x) dx \right| \\ &\lesssim \sup_{\|g\|_{L^2(B)} \leq 1} \sum_{k=0}^{\infty} \exp \left\{ -\frac{2^k r_B^2}{c r_B^2} \right\} \|f - f_B\|_{L^2(U_k(B))} \|g\|_{L^2(B)} \end{aligned}$$

$$\begin{aligned}
&\lesssim \sum_{k=0}^{\infty} e^{-2^k/c} \left\{ \|f - f_{2^k B}\|_{L^2(U_k(B))} + |U_k(B)|^{1/2} |f_B - f_{2^k B}| \right\} \\
&\lesssim \sum_{k=0}^{\infty} e^{-2^k/c} 2^{kn/2} \|f - f_{2^k B}\|_{L^2(U_k(B))},
\end{aligned}$$

where c is a positive constant. Since $f \in C_c(\mathbb{R}^n) \subset \text{CMO}(\mathbb{R}^n)$, by the dominated convergence theorem for series, [10, Proposition 3.5] and the fact that $\text{CMO}(\mathbb{R}^n) \subset \text{VMO}(\mathbb{R}^n)$, we have

$$\begin{aligned}
\gamma_1(f) &= \lim_{c \rightarrow 0} \sup_{B: r_B \leq c} \frac{1}{|B|^{1/2}} \|(I - e^{-r_B^2 L})^M f\|_{L^2(B)} \\
&\lesssim \sum_{k=0}^{\infty} e^{-2^k/c} 2^{kn} \lim_{c \rightarrow 0} \sup_{B: r_B \leq c} \frac{1}{|2^k B|^{1/2}} \|f - f_{2^k B}\|_{L^2(U_k(B))} = 0.
\end{aligned}$$

Similarly, we have that $\gamma_2(f) = \gamma_3(f) = 0$, which implies that $f \in \text{VMO}_L(\mathbb{R}^n)$, and hence $C_c(\mathbb{R}^n) \subset \text{VMO}_L(\mathbb{R}^n)$. Since $C_c(\mathbb{R}^n)$ is dense in $\text{CMO}(\mathbb{R}^n)$, by the fact that for all $g \in \text{CMO}(\mathbb{R}^n)$, $\|g\|_{\text{VMO}_L(\mathbb{R}^n)} = \|g\|_{\text{BMO}_L(\mathbb{R}^n)} \lesssim \|g\|_{\text{BMO}(\mathbb{R}^n)} \sim \|g\|_{\text{CMO}(\mathbb{R}^n)}$, we finally obtain that $\text{CMO}(\mathbb{R}^n) \subset \text{VMO}_L(\mathbb{R}^n)$.

On the other hand, if $n \geq 3$, then there exist L as in (1.2) and $p \in (1, 2)$ such that $\nabla(L^*)^{-1/2}$ is not bounded on $L^p(\mathbb{R}^n)$; see [1, 14]. By the fact that the Riesz transform $\nabla(L^*)^{-1/2}$ is bounded on $L^2(\mathbb{R}^n)$ and from $H_{L^*}^1(\mathbb{R}^n)$ to $L^1(\mathbb{R}^n)$ (see [16, 20]), we obtain that $H_{L^*}^1(\mathbb{R}^n)$ is a proper subspace of $H^1(\mathbb{R}^n)$. Otherwise, $H_{L^*}^1(\mathbb{R}^n) = H^1(\mathbb{R}^n)$, then by the interpolation theorem, $\nabla(L^*)^{-1/2}$ is bounded on $L^p(\mathbb{R}^n)$ for all $p \in (1, 2)$, which contradicts with the fact that $\nabla(L^*)^{-1/2}$ is not bounded on $L^p(\mathbb{R}^n)$ for some $p \in (1, 2)$.

Now by Theorem 4.2, we have that

$$(\text{VMO}_L(\mathbb{R}^n))^* = H_{L^*}^1(\mathbb{R}^n) \subsetneq H^1(\mathbb{R}^n) = (\text{CMO}(\mathbb{R}^n))^*,$$

which implies that $\text{CMO}(\mathbb{R}^n) \subsetneq \text{VMO}_L(\mathbb{R}^n)$, and hence, completes the proof of Proposition 4.2. $\square$

From Proposition 4.2, we deduce the following conclusion.

Corollary 4.1. *When A has real entries, or when the dimension $n = 1$ or $n = 2$ in the case of complex entries, the spaces $\text{CMO}(\mathbb{R}^n)$ and $\text{VMO}_L(\mathbb{R}^n)$ coincide with equivalent norms.*

To prove Corollary 4.1, we need the following lemma. (We are very grateful to the referee for him/her to tell us Corollary 4.1 and its proof.)

Lemma 4.5. *Let X and Y be two Banach spaces such that $X \subset Y$ with continuous embedding. Assume that X^* and Y^* coincide with equivalent norms. Then $X = Y$ with equivalent norms.*

Proof. Let $J : X \rightarrow Y$ be the inclusion map, which is a continuous linear map by assumption. As is easily seen, its adjoint $J^* : Y^* \rightarrow X^*$ is the map which, to any $\phi \in Y^*$, associates its restriction to X . The Hahn-Banach theorem yields that J^* is onto. Moreover, it is also one-to-one. Indeed, if this was not true, there would exist $\phi \in Y^*$ identically vanishing on X but not on Y , which would contradict the assumption that X^* and Y^* coincide with equivalent norms. Thus, J^* is a continuous isomorphism, and, by Theorem 4.15 in [23], J is an isomorphism between X and Y , which exactly means that $X = Y$. From this and Corollaries 2.12(c) of the open mapping theorem in [23, pp. 49-50], it follows that X and Y coincide with equivalent norms, which completes the proof of Lemma 4.5. $\square$

Proof of Corollary 4.1. When A has real entries, or when the dimension $n = 1$ or $n = 2$ in the case of complex entries, the heat kernel always satisfies the Gaussian pointwise estimates in size and regularity (see [6]), and the spaces $H_{L^*}^1(\mathbb{R}^n)$ and $H^1(\mathbb{R}^n)$ coincide with equivalent norms (see [5, 27]). By this and Proposition 4.2, we obtain that $\text{CMO}(\mathbb{R}^n) \subset \text{VMO}_L(\mathbb{R}^n)$ and $(\text{CMO}(\mathbb{R}^n))^* = H^1(\mathbb{R}^n) = H_{L^*}^1(\mathbb{R}^n) = (\text{VMO}_L(\mathbb{R}^n))^*$, which together with Lemma 4.5 imply that $\text{CMO}(\mathbb{R}^n)$ and $\text{VMO}_L(\mathbb{R}^n)$ coincide with equivalent norms. This finishes the proof of Corollary 4.1. $\square$

Acknowledgment

The authors sincerely wish to express their deeply thanks to the referee for her/his very carefully reading and also her/his so many valuable and helpful remarks which made this article more readable. In particular, Corollary 4.1 and its proof belong to the referee.

References

- [1] P. Auscher, On necessary and sufficient conditions for L^p -estimates of Riesz transforms associated to elliptic operators on $\mathbb{R}^n$ and related estimates, *Mem. Amer. Math. Soc.* 186 (2007), 1-75.
- [2] P. Auscher, X. T. Duong and A. McIntosh, Boundedness of Banach space valued singular integral operators and Hardy spaces, *Unpublished Manuscript*, 2005.
- [3] P. Auscher, S. Hofmann, M. Lacey, A. McIntosh and Ph. Tchamitchian, The solution of the Kato square root problem for second order elliptic operators on $\mathbb{R}^n$, *Ann. of Math.* (2) 156 (2002), 633-654.
- [4] P. Auscher, A. McIntosh and E. Russ, Hardy spaces of differential forms on Riemannian manifolds, *J. Geom. Anal.* 18 (2008), 192-248.
- [5] P. Auscher and E. Russ, Hardy spaces and divergence operators on strongly Lipschitz domains of $\mathbb{R}^n$, *J. Funct. Anal.* 201 (2003), 148-184.
- [6] P. Auscher and P. Tchamitchian, Square root problem for divergence operators and related topics, *Astérisque*, 249 (1998), 1-172.
- [7] G. Bourdaud, Remarques sur certains sous-espaces de $\text{BMO}(\mathbb{R}^n)$ et de $\text{bmo}(\mathbb{R}^n)$, *Ann. Inst. Fourier (Grenoble)* 52 (2002), 1187-1218.
- [8] R. R. Coifman, Y. Meyer and E. M. Stein, Some new functions and their applications to harmonic analysis, *J. Funct. Anal.* 62 (1985), 304-335.

- [9] R. R. Coifman and G. Weiss, Extensions of Hardy spaces and their use in analysis, *Bull. Amer. Math. Soc.* 83 (1977), 569-645.
- [10] D. Deng, X. T. Duong, L. Song, C. Tan and L. Yan, Functions of vanishing mean oscillation associated with operators and applications, *Michigan Math. J.* 56 (2008), 529-550.
- [11] X. T. Duong and L. Yan, New function spaces of BMO type, the John-Nirenberg inequality, interpolation, and applications, *Comm. Pure Appl. Math.* 58 (2005), 1375-1420.
- [12] X. T. Duong and L. Yan, Duality of Hardy and BMO spaces associated with operators with heat kernel bounds, *J. Amer. Math. Soc.* 18 (2005), 943-973.
- [13] C. Fefferman and E. M. Stein, H^p spaces of several variables, *Acta Math.* 129 (1972), 137-193.
- [14] J. Frehse, An irregular complex valued solution to a scalar uniformly elliptic equation, *Calc. Var. Partial Differential Equations* 33 (2008), 263-266.
- [15] S. Hofmann and J. M. Martell, L^p bounds for Riesz transforms and square roots associated to second order elliptic operators, *Publ. Mat.* 47 (2003), 497-515.
- [16] S. Hofmann and S. Mayboroda, Hardy and BMO spaces associated to divergence form elliptic operators, *Math. Ann.* 344 (2009), 37-116.
- [17] S. Hofmann and S. Mayboroda, Correction to "Hardy and BMO spaces associated to divergence form elliptic operators", arXiv: 0907.0129.
- [18] S. Janson, Generalizations of Lipschitz spaces and an application to Hardy spaces and bounded mean oscillation, *Duke Math. J.* 47 (1980), 959-982.
- [19] R. Jiang and D. Yang, Predual spaces of Banach completions of Orlicz-Hardy spaces associated with operators, arXiv: 0906.1880.
- [20] R. Jiang and D. Yang, New Orlicz-Hardy spaces associated with divergence form elliptic operators, *J. Funct. Anal.* (to appear).
- [21] R. Jiang, D. Yang and Y. Zhou, Orlicz-Hardy spaces associated with operators, *Sci. China Ser. A*, 52 (2009), 1042-1080.
- [22] F. John and L. Nirenberg, On functions of bounded mean oscillation, *Comm. Pure Appl. Math.* 14 (1961), 415-426.
- [23] W. Rudin, *Functional Analysis*, Second Edition, McGraw-Hill, Inc., New York, 1991.
- [24] D. Sarason, Functions of vanishing mean oscillation, *Trans. Amer. Math. Soc.* 207 (1975), 391-405.
- [25] B. E. Viviani, An atomic decomposition of the predual of $BMO(\rho)$, *Rev. Mat. Iberoamericana* 3 (1987), 401-425.
- [26] W. Wang, The predual spaces of tent spaces and some characterizations of $\lambda_\alpha(\mathbb{R}^n)$ spaces, *Beijing Daxue Xuebao* 24 (1988), 535-551.
- [27] L. Yan, Classes of Hardy spaces associated with operators, duality theorem and applications, *Trans. Amer. Math. Soc.* 360 (2008), 4383-4408.
- [28] K. Yosida, *Functional Analysis*, Sixth Edition, Springer-Verlag, Berlin, 1978.

Renjin Jiang:

School of Mathematical Sciences, Beijing Normal University

Laboratory of Mathematics and Complex Systems, Ministry of Education

100875 Beijing

People's Republic of China

E-mail: `rj-jiang@mail.bnu.edu.cn`

Present address

Department of Mathematics and Statistics, University of Jyväskylä

P.O. Box 35 (MaD), 40014, Finland

Dachun Yang (corresponding author)

School of Mathematical Sciences, Beijing Normal University

Laboratory of Mathematics and Complex Systems, Ministry of Education

100875 Beijing

People's Republic of China

E-mail: `dcyang@bnu.edu.cn`